


“Quantum and classical chaos: What comes next?”

Symposium in honor of
Prof. Marko Robnik
on the occasion of his 60th birthday

Organized by the
Faculty of Mathematics and Physics, University of Ljubljana
Ljubljana 9-11 October 2014



Forward

Nonlinear dynamics has been a vivid area of multidisciplinary research for several past decades with numerous successful applications in diverse areas of science. From a physicist's perspective, one is still intrigued by the connection between the classical (often completely deterministic) chaotic dynamics and the properties of the corresponding quantum counterpart.

The key aspect which often unifies such connections is universality. The aim of the Symposium is to bring together renowned scientists from different fields, sharing a common interest in complex classical and quantum systems, with a particular emphasis on discussing open problems and perspectives in classical and quantum dynamics, from viewpoints of fundamental research and applications.

The venue of the symposium as well as lodging of all participants is taking place at the Grand Hotel Union, Ljubljana.

Organizing committee

Prof. Tomaz Prosen, University of Ljubljana, Slovenia

Dr. Thanos Manos, Research Center Jülich, Germany

Prof. Gregor Veble, Pipistrel and University of Nova Gorica, Slovenia



Professor Marko Robnik

The scientific work of Professor Marko Robnik is in the wide domain of theoretical physics, and it began in 1978 in the field of mathematical physics, kinetic theory and fluid mechanics at the University of Ljubljana, followed by theoretical astrophysics at the University of Bonn, mainly treating physical processes in astrophysics, and nonlinear dynamics of classical and quantum chaos, later also at the University of Bristol, England, Max-Planck-Institute for Nuclear Physics in Heidelberg, Germany, and Institute of Theoretical Physics, University of California at Santa Barbara, USA, in 1989. Since then he is working mainly on classical and quantum chaos, and mathematical physics. Marko Robnik is best known for his seminal work, together with M. Berry, on energy level spacing distribution in quantized Hamiltonian systems with generic, mixed phase space where regular and chaotic motion can coexist. In another seminal paper he clarified the role of anti-unitary symmetry in statistics of energy levels, and proposed popular Aharonov-Bohm quantum billiard models. A particularly simple and appealing billiard model whose shape is given as a conformal quadratic mapping of a unit disk is often known as Robnik billiard. He is the founder of the CAMTP - Center for Applied Mathematics and Theoretical Physics, at the University of Maribor, in 1990.

CAMTP was founded when he returned to Slovenia from the Institute of Theoretical Physics, University of California at Santa Barbara, USA. With the great moral and financial support of the Slovenian Government on the one hand, and the operational support by the Rector Professor Alojz Križman at the time on the other hand, the Center started its scientific research activities in September 1990. It became an independent research institution and legal entity in July 1991, but still academically part of the University of Maribor. Professor Marko Robnik is the director of the institute since the foundation.

The main purpose of the Center is to carry out research in theoretical physics and applied mathematics, mainly in nonlinear dynamics, and also to educate young researchers, PhD students, and also to organize scientific meetings, mainly international, but also elite national meetings with strong international participation. The Center is small by the number of members but nevertheless very active, at the moment (January 2014) it has nine members, from four countries.

So far at CAMTP five PhD students of theoretical physics have successfully finished their PhD thesis work, among them 3 Slovenians, one from Greece and one from Brazil. One Slovenian earned PhD in mathematics. Two more PhD students of physics are working at CAMTP towards earning the PhD degree, and one in mathematics.

One should emphasize the strong international conference activities of the Center, as it is organizing 2-3

international and high level or even world top level conferences per year. Since 1993 the Center has organized more than 51 scientific conferences and meetings, namely seven series of scientific meetings:

i. *The international summer schools and conferences Let's Face Chaos through Nonlinear Dynamics*. They started them in 1993, and had 9 so far, every three years, in Maribor, with two working weeks, with about 80-100 participants each time. The last one was on 22 June - 6 July 2014.

ii. *Japan-Slovenia Seminars on Nonlinear Science*. They meet almost every year in Maribor or in Japan, have typically 3 working days in Maribor, and twice three days in Japan, at various top universities, in Tokyo, Kyoto, Nara and Osaka. Typically about 40-50 participants. So far there were 13 meetings. The next one will be in October 2014 in Tokyo and Kansai area (Osaka, Kyoto, Nara).

iii. *European Advanced Studies Conferences*, organized in collaboration with Technical University of Munich, Germany, and the University of Patras, Greece, almost every year, with 3-4 working days. They had 8 so far. They take place at various places in Europe.

iv. *SOCRATES Workshops on nonlinear dynamics*, so far they had five, twice in Marburg, Germany, twice in Maribor, Slovenia, in collaboration with Professor Hans-Jürgen Stöckmann of University of Marburg. (Later there was one follow-up meeting in Freiburg, in Germany, independent of CAMTP.) Each time about 30 participants for one week, typically.

v. *Christmas Symposia on Physics*, every year since 2002, so far 12 meetings, the next one will be in December of 2014. There are three working days and about 45 participants.

vi. *Computer Algebra and Scientific Computing*, so far three meetings (2010, 2012 and 2013), about 45 participants each time and three to four working days.

vii. *1st Out of the Box Conference*, in collaboration with the University of Maribor and Rector Prof. Dr. Danijel Rebolj, 15-17 May 2012, with the subtitle "Innovative Ways to Improve the Culture of Living", with 30 invited speakers, including three Nobel Prize Laureates, and 27 other eminent speakers, among them Prof. Dr. Dr. h. c. mult. Felix Unger, President of the EASA, Prof. Dr. Dr. h. c. mult. Siegfried Großmann from the University of Marburg, Germany, etc.

Invited speakers

1. Bäcker Arnd (Technical University Dresden, Germany)
2. Bountis Tassos (University of Patras, Greece)
3. Casati Giulio (University Insubria, Como, Italy)
4. Cvetič Mirjam (University of Pennsylvania, USA, and CAMTP, University of Maribor, Slovenia)
5. Dvorak Rudolf (University of Vienna, Austria)
6. Flach Sergej (Massey University, Auckland, New Zealand)
7. Geisel Theo (Max Planck Institute for Dynamics and Self-Organization, Göttingen, Germany)
8. Haake Fritz (University Duisburg-Essen, Germany)
9. Kottos Tsampikos (Wesleyan University, USA)
10. Krylov George (Belarus State University, Minsk, Belarus)
11. Kuvshinov Viatcheslav (Joint Institute for Power and Nuclear Research, Minsk, Belarus)
12. Manos Thanos (Research Center Jülich, Germany)
13. McClintock Peter V. E. (Lancaster University, UK)
14. Plessas Willibald (University of Graz, Austria)
15. Richter Klaus (University of Regensburg, Germany)
16. Robnik Marko (CAMTP, University of Maribor, Slovenia)
17. Saksida Pavle (University of Ljubljana, Slovenia)
18. Seligman Thomas (UNAM, Mexico)
19. Stefanovska Aneta (Lancaster University, UK)
20. Stöckmann Hans-Juergen (Philipps-Universität Marburg, Germany)
21. Takatsuka Kazuo (University of Tokyo, Japan)
22. Weidenmüller Hans A. (Heidelberg University, Germany)
23. Žnidarič Marko (University of Ljubljana, Slovenia)

Program

Wednesday, 8.10.2014

19:30 — welcome dinner —

Thursday, 9.10.2014	Friday, 10.10.2014	Saturday, 11.10.2014
Chair: Bountis	Chair: Stöckmann	Chair: Plessas
09:00 Welcome	09:00 Haake	09:00 Stöckmann
09:15 Plessas	09:45 Casati	09:45 Kottos
10:00 Stefanovska	10:30 Seligman	10:30 Flach
10:45 Dvorak		
11:30 -coffee break, discussions-	11:15 -coffee break, discussions-	11:15 -coffee break, discussions-
12:00 Kuvshinov	11:45 Robnik	11:45 Bountis
12:45 McClintock	12:30 Richter	12:30 Žnidarič
13:30 — lunch —	13:15 — lunch —	13:15 — lunch —
Chair: Casati	Chair: McClintock	
15:30 Krylov	15:15 Cvetič	
16:15 Manos	16:00 Weidenmueller	
17:00 -coffee break, discussions-	16:45 -coffee break, discussions-	
17:30 Takatsuka	17:15 Bäcker	
18:15 Saksida	18:00 Geisel	
19:00 — discussions —	19:00 — Reception & concert —	
19:30 — dinner —	20:30 — Festive dinner —	

Abstracts of Talks

Visualization of the hierarchy of regular tori in 4d maps

Arnd Bäcker

*Technische Universität Dresden, Institut für Theoretische Physik and
Center for Dynamics, 01062 Dresden, Germany
Max-Planck-Institut für Physik komplexer Systeme,
Nöthnitzer Straße 38, 01187 Dresden, Germany*

For a generic 4D symplectic map we visualize the global organization of regular tori using 3D phase-space slices. The regular 2-tori are shown to be arranged around a skeleton of elliptic 1-tori. The 1-tori occur in two types of one-parameter families: (so-called) Lyapunov families emanating from elliptic-elliptic periodic orbits and families originating from rank-1 resonances. At resonance gaps of both types of families periodic orbits may occur, similar to the Poincaré-Birkhoff theorem for 2D maps. Based on this we describe the hierarchical structure of regular tori in the 4D phase space. Each level of the hierarchy can result from three types of structures which generalizes the islands-around-islands hierarchy in 2D maps. As an application we consider the representation of eigenstates of the corresponding quantum map to investigate the semiclassical eigenfunction hypothesis.

References

- [1] S. Lange, M. Richter, F. Onken, A. Bäcker, and R. Ketzmerick, Global structure of regular tori in a generic 4D symplectic map, *Chaos* 24, 024409 (2014)
- [2] M. Richter, S. Lange, A. Bäcker, and R. Ketzmerick, Visualizing classical structures and quantum states of 4D maps by 3D phase-space slices, *Phys. Rev. E* 89, 022902 (2014).
- [3] For videos of 3D phase space slices see: <http://www.comp-phys.tu-dresden.de/supp/>

Complexity Aspects of Hamiltonian Dynamics and Statistics

Tassos Bountis

*Department of Mathematics and
Center for Research and Applications of Nonlinear Systems,
University of Patras, Patras 26500 Greece*

As is well known, many problems of classical mechanics, celestial mechanics and solid state physics, when dissipation is neglected, are expressed in Hamiltonian form and deal with multi-dimensional systems of interacting mass particles. These typically execute oscillatory motions near stable equilibria and the most frequently asked questions concern the amplitudes and frequencies of these oscillations. However, since the interactions are fundamentally nonlinear, unstable equilibria become important as the total energy increases, causing the motion to drift into chaotic domains where the oscillations become unpredictable and statistical mechanics is expected to take over. In this presentation, I will first mention some very efficient techniques for distinguishing regular from chaotic domains in Hamiltonian systems of many degrees of freedom. More importantly, I will describe some complex aspects of Hamiltonian dynamics and statistics, demonstrating “hierarchies” of order, weak and strong chaos and discuss the role of long vs. short range interactions.

Dedication: *This lecture is dedicated to Professor Marko Robnik, with affection for his 60th birthday and appreciation for his contributions to the field of Hamiltonian systems, classical and quantum.*

References

1. T. Bountis and Haris Skokos, “Complex Hamiltonian Dynamics”, Synergetics series of Springer Verlag, April 2012.
2. C. Antonopoulos, T. Bountis and V. Basios, “Quasi--Stationary Chaotic States in Multi-Dimensional Hamiltonian Systems”, Physica A, Vol. **390**, 3290-3307. (2011).
3. H. Christodoulidi, C. Efthymiopoulos and T. Bountis, “Energy Localization on q-Tori, Long Term Stability and the Interpretation of the FPU Paradox”, PRE **81**, 016210 (2010).
4. Ch. Skokos, T. Bountis and C. Antonopoulos, “Geometrical Properties of Local Dynamics in Hamiltonian Systems: The Generalized Alignment (GALI) Method”, Physica D **231**, 30 (2007).

Conservation laws, symmetry breaking and control of the heat current

Giulio Casati

*Center for Complex System University of Insubria
and
Lake Como School of Advanced Studies
Como - Italy*

The understanding of the underlying dynamical mechanisms which determines the macroscopic laws of heat conduction is a long standing task of non-equilibrium statistical mechanics. A better understanding of such mechanism may also lead to potentially interesting applications based on the possibility to control the heat flow. Indeed, a model of thermal rectifier has been recently proposed in which heat can flow preferentially in one direction. The proposed rectifying mechanism is of very general nature and theoretical and experimental work is currently undergoing.

In particular we discuss here the possibility to increase the rectifying factor by orders of magnitude. We then discuss a new approach, which is rooted in nonlinear dynamical systems, for increasing the efficiency of thermoelectric machines. The main focus will be on the physical mechanisms, unveiled by these dynamical models, which lead to high thermoelectric efficiency, approaching the Carnot limit. The role of conservation laws and symmetry breaking will be emphasized.

String Theory Confronts Particle Physics and Black Holes

Mirjam Cvetič

University of Pennsylvania, USA, and CAMTP, University of Maribor, Slovenia

String theory, as the prime candidate for quantum unification of particle physics and gravity, sheds light on important fundamental questions such as the microscopic structure of black holes and the geometric origin of particle physics. We review these developments such as the introduction of extended objects - Dirichlet branes - and highlight the important geometric role of these objects in deriving particle physics from string theory, including recent progress within string theory at finite coupling (F-theory), as well as their role in deriving the black hole microscopics, including recent progress for non-extremal black holes.

Planetary systems and the formation of habitable planets

Rudolf Dvorak

University of Vienna, Austria

Since the last 20 years many planets around other system were detected either from the ground or by space missions. More than 1100 planetary systems with more than one discovered planet are known but for planets being habitable many conditions need to be fulfilled: e.g. the spectral type and age of the hosting star, the location in our galaxy and also the orbit of a planet and its physical nature. Different estimates lead to quite different results especially for planet which host in fact life. In this talk different results and aspects concerning this question will be addressed.

Quantum chaotic subdiffusion in random potentials

M.V. Ivanchenko¹, T.V. Laptyeva² and **S. Flach**³

¹*Department for Bioinformatics, Lobachevsky State University of Nizhniy Novgorod, Russia*


²*Theory of Control Department, Lobachevsky State University of Nizhniy Novgorod, Russia*

³*New Zealand Institute for Advanced Study, Massey University, Auckland, New Zealand*

Two interacting particles (TIP) in a disordered chain propagate beyond the single particle localization length ξ_1 up to a scale $\xi_2 > \xi_1$. An initially strongly localized TIP state expands almost ballistically up to ξ_1 . The expansion of the TIP wave function beyond the distance $\xi_1 \gg 1$ is governed by highly connected Fock states in the space of noninteracting eigenfunctions. The resulting dynamics is subdiffusive, and the second moment grows as $m_2 \sim t^{1/2}$ [1], precisely as in the strong chaos regime for corresponding nonlinear wave equations [2-7]. This surprising outcome stems from the huge Fock connectivity and resulting quantum chaos. The TIP expansion finally slows down towards a complete halt - in contrast to the nonlinear case.

References

- [1] M. V. Ivanchenko, T. V. Laptyeva and S. Flach, *Phys.Rev. B Rapid Comm.* **89**, 060301 (2014).
- [2] S. Flach, D. O. Krimer and Ch. Skokos, *Phys. Rev. Lett.* **102**, 024101 (2009).
- [3] S. Flach, *Chem. Phys.* **375**, 548 (2010).
- [4] T.V. Laptyeva et al, *EPL* **91**, 30001 (2010).
- [5] J.D. Bodyfelt et al, *Phys. Rev. E*, **84**, 016205 (2011).
- [6] S. Flach, arXiv:1405.1122 (2014).
- [7] T. V. Laptyeva, M. V. Ivanchenko and S. Flach, arXiv:1407.1441 (2014).



Rhythms, Algorithms, and Self-Similarity in Music

Theo Geisel

*Max Planck Institute for Dynamics and Self-Organization &
Bernstein Center for Computational Neuroscience Göttingen, Germany*

Even the best musicians do not play rhythms with perfect precision. Slight deviations from an ideal beat pattern are a fundamental characteristic of music played by humans. This talk investigates the statistical laws underlying rhythmic fluctuations and their role in musical perception. Based on these findings one can make computer generated music sound more human. Audio examples from the Art of Fugue to stochastic music highlight the general role of long range correlations and self-similarity in music for its perception by the information processing in our brains.

Different paths to spectral universality

Fritz Haake

Universität Duisburg-Essen

Numerical case studies reveal that dynamics with full chaos in their classical limit show universal fluctuations in their two-point function of the quantum level density, at least within the window(s) of (quasi-)energies where correlations persist above some small noise level. Good understanding of such universality has by now been reached by four different methods:

- (i) Random matrix theory provides ensembles of (Hermitian and unitary) matrices phenomenologically describing symmetry classes of Hamiltonians of Floquet maps and predicts closed-form expressions for the two-point function within certain symmetry classes; moreover, the variances of that function are shown to be inversely proportional to the matrix size N such that for $N \rightarrow \infty$ all matrices have the same (universal) spectral fluctuations.
- (ii) 'Level dynamics' maps the dependence of quantum levels on some control parameter to the time evolution of some fictitious classical many-particle system whose equilibrium statistics is equivalent to random-matrix theory.
- (iii) Gutzwiller's periodic-orbit theory allows to calculate spectral characteristics like the two-point function as the sum of contributions from bunches of near action degenerate orbits, thus recovering the RMT results for individual quantum systems.
- (iv) The field theoretic method known as supersymmetric sigma model reveals the periodic-orbit sums as perturbation expansions and even comes with more powerful non-perturbative procedures.

I shall briefly describe the relative status of the four approaches.

Random Matrix Theory approach to Thermal Transport in Complex Lattices

Tsampikos Kottos

Wesleyan University, USA

Using random matrix theory modeling we investigate the thermal transport in networks of N oscillators coupled harmonically with random springs. When the network is assumed to be fully connected we show, that both the average phonon heat current and its variance are scale invariant and take universal values in the large N limit. These anomalous mesoscopic fluctuations are the hallmark of strong correlations between normal modes. In the case of quasi-one dimensional networks with finite connectivity between nearby oscillators, we show that the heat current obeys a one-parameter scaling law. The resulting β function indicates that an anomalous Fourier law is applicable in the diffusive regime, while in the localization regime the heat current decays exponentially with the sample size. Our approach opens a new way to investigate the effects of Anderson localization in heat conduction based on the powerful ideas of scaling theory and random matrix theory modeling.

Charge transport in single layer graphene: beyond massless Dirac fermion approach

Halina Grushevskaya, George Krylov and Maxim Belov

Belarus State University, Minsk, Belarus

Approaches to theoretical description of charge transport in single layer graphene are discussed. Consequent quantum field theory approach has been proposed. It generalizes known massless Dirac fermion model and is based on the Dirac – Hartree – Fock self-consistent field approximation and assumption on antiferromagnetic ordering of graphene lattice. The developed approach allows asymmetric charged carriers in single layer graphene with partially degenerated Dirac cones. Some open problems of graphene physics are also discussed.

Chaotic and Unstable Behaviour of Quantum Colour Particles in QCD Vacuum

Vyacheslav Kuvshinov

Joint Institute for Power and Nuclear Research, Minsk, Belarus

Quantum colour particles are considered in stochastic vacuum of Quantum Chromodynamics (QCD) as environment, with density matrix averaged over external QCD vacuum degrees of freedom. Density matrices decoherence rates, purities, Von Neumann entropies, fidelities are calculated for different pure, mixed, separable, entangled colour states. It is shown that in confinement region (Wilson loop decays exponentially) colour particle movement is chaotic and unstable, order-chaos transition appears with critical energy depended on Higgs mass and coupling constant, density matrixes lose nondiagonal elements, final states become mixed and colourless, quantum squeezing and entanglement of colour particles appear.

Probing the role of accelerator modes on the dynamical localization properties of the quantum kicked rotator and on the anomalous diffusion of its classical analogue

Thanos Manos¹ and Marko ROBNIK²

¹*Institute of Neuroscience and Medicine, Research Centre Jülich, Germany*

²*CAMTP-Center for Applied Mathematics and Theoretical Physics, University of Maribor, Slovenia*

In the first part of the talk, we study the N -dimensional model of the quantum kicked rotator in the classically fully chaotic regime, which in the limit of sufficiently large N tends to the quantized kicked rotator. We describe the features of dynamical localization of chaotic eigenstates as a paradigm for other both time-periodic and time-independent (autonomous) fully chaotic or/and mixed type Hamilton systems. We generalize the scaling variable to the case of anomalous diffusion in the classical phase space, by deriving the localization length for the case of generalized classical diffusion. We greatly improve the accuracy and statistical significance of the numerical calculations, giving rise to the following conclusions: (i) The level spacing distribution of the eigenphases is very well described by the Brody distribution, systematically better than by other proposed models. (ii) We study the eigenfunctions of the Floquet operator and characterize their localization properties using the information entropy measure. (iii) We show the existence of a scaling law between the localization parameter and the relative localization length, now including the regimes of anomalous diffusion.

In the second part, we focus on the effect of the anomalous diffusion arising due to the accelerator modes in the classical kicked rotator, exemplified by the standard map. The systematic approach rests upon detecting the regular and chaotic regions and thus to describe in detail the structure of the phase space, the description of the momentum distribution in terms of the Lévy stable distributions, the numerical calculation of the diffusion exponent and of the corresponding diffusion constant. We use this approach to analyze in detail and systematically the standard map at all values of the kick parameter K , up to $K=70$. All complex features of the anomalous diffusion are well understood in terms of the role of the accelerator modes, mainly of period 1 at large $K \geq 2\pi$, but also of higher periods (2,3,4,...) at smaller values of $K \leq 2\pi$.

References

- [1] Izrailev F M 1988 *Phys. Lett. A* **134** 13
- [2] Izrailev F M 1990 *Phys. Rep.* **196** 299
- [3] Batistić B and Robnik M 2010 *J. Phys. A: Math. Theor.* **43** 215101
- [4] Manos T and Robnik M 2013 *Phys. Rev. E* **87** 062905
- [5] Batistić B, Manos T and Robnik M 2013 *EPL* **102** 50008
- [6] Batistić B and Robnik M 2013 *J. Phys. A: Math. Theor.* **46** 315102
- [7] Manos T and Robnik M 2014 *Phys. Rev. E* **89** 022905

Coulomb blockade dynamics in biological ion channels

William A T Gibby,¹ Igor Kh Kaufman,¹ Dmitri G Luchinsky,^{1,2} **Peter V E McClintock**¹
and Robert S (Bob) Eisenberg³

¹*Department of Physics, Lancaster University, Lancaster LA1 4YB, UK*

²*Mission Critical Technologies Inc., 2041 Rosecrans Ave. Suite 225 El Segundo, CA 90245, USA*

³*Department of Molecular Biophysics and Physiology, Rush Medical College, 1750 West Harrison, Chicago, IL 60612, USA*

Biological ion channels are natural nanopores that provide for fast and selective permeation of physiologically important ions (e.g. Na⁺, K⁺ and Ca²⁺) through cellular membranes. Although a great deal is now known about ion channels, including their detailed atomic structure in some cases, this information has helped surprisingly little in understanding how they actually work: the central conundrum – their ability to conduct almost at the rate of free diffusion while remaining highly selective for particular ions – remains largely unresolved. We show that treating it as a problem in nonlinear dynamics under the influence of electrostatic forces can account for numerous experimental observations.

In particular, we hypothesise that many of the complex permeation and selectivity phenomena of biological ion channels can be described in terms of a much simpler and well-studied system – the quantum dot.¹ We infer that permeation is governed by ionic Coulomb blockade (ICB),² a discrete electrostatic phenomenon closely analogous to its quantal counterpart.¹ It immediately explains the conduction bands seen in Brownian dynamics simulations³ as the fixed negative charge Qf at the selectivity filter (SF) is varied. They are Coulomb blockade oscillations⁴ between stop bands Z_n (blockade by an ion trapped in the channel, cf. an electron trapped in a dot) and conduction bands M_n corresponding to resonant conduction. Singular points of the ICB oscillations are given by

$$\begin{aligned} Z_n &= zen \pm \delta Z_n, & \text{Coulomb blockade} \\ M_n &= ze(n + 1/2) \pm \delta M_n & \text{Resonant conduction} \end{aligned} \quad (1)$$

where e is the elementary charge, z the valence, $n = 0, 1, 2, \dots$ the number of ions captured at the SF, and $\delta Z_n, \delta M_n$ are possible band shifts to account for affinity, ion-ion interactions and hydration.

Equation (1) is similar to its electronic counterpart¹ but the positions M_n, Z_n of the ICB bands are proportional to z , thereby providing an explanation for valence selectivity. The model also enables bands to be identified with particular channels⁵ and accounts for the shapes of the conduction bands and occupancy transitions.^{3,6} The detailed geometry of the SF has little influence on valence selectivity, for which the magnitude of Qf is the dominant factor, but it may be very important for alike charge selectivity: the explicit inclusion of hydration effects⁷ in Eqs. (1) promises to pave the way to the first generalised model of selectivity. The specific prediction of the ICB model (1) – the existence of a universal periodic pattern of conduction bands M_n separated by stop bands Z_n – is already well-supported by simulations and by the identification of mutant channels.^{3,8}

References

1. Alhassid, Y., Statistical theory of quantum dots, *Rev. Mod. Phys.* **72**, 895–968 (2000).

2. Krems, M. & Di Ventra, M., Ionic coulomb blockade in nanopores, *J. Phys. Condens. Matter* **25**, 065101 (2013).
3. Kaufman, I. K., Tindjong, R., Luchinsky, D. G., McClintock, P. V. E. & Eisenberg, R. S., Resonant multi-ion conduction in a simple model of calcium channels, in *22nd Intern. Conf. on Noise and Fluctuations (ICNF)*, Montpellier, 24–28 June 2013 (Eds. J. M. Routoure, L. Varani & F. Pascal), p. doi: 10.1109/ICNF.2013.6578926 (IEEE Conf. Proc., 2013).
4. Beenakker, C. W. J., Theory of Coulomb-blockade oscillations in the conductance of a quantum dot, *Phys. Rev. B* **44**, 1646–1656 (1991).
5. Kaufman, I. K., Luchinsky, D. G., Tindjong, R., McClintock, P. V. E. & Eisenberg, R. S., Energetics of discrete selectivity bands and mutation-induced transitions in the calcium-sodium ion channels family, *Phys. Rev. E* **88**, 052712 (2013).
6. Kaufman, I. K., McClintock, P. V. E. & Eisenberg, R. S., Ionic Coulomb blockade and resonant conduction in biological ion channels, *arXiv*: 1405.1391 (2014).
7. Eisenman, G. & Horn, R., Ionic selectivity revisited: the role of kinetic and equilibrium processes in ion permeation through channels, *J. Membrane Biol.* **76**, 197–225 (1983).
8. Kaufman, I. K., Luchinsky, D. G., Tindjong, R., McClintock, P. V. E. & Eisenberg, R. S., Multi-ion conduction bands in a simple model of calcium ion channels, *Phys. Biol.* **10**, 026007 (2013).

Quantum Chaos in Hadron Spectra

Wilibald Plessas

*Theoretical Physics, Institute of Physics, University of Graz
Universitaetsplatz 5, 8010 Graz, Austria*

Quantum-chaotic behaviour has been identified in quantum spectra quite some time ago according to the Bohigas-Giannoni-Schmit conjecture using random matrix theory [1]. We shall discuss the appearance of quantum chaos in relativistically invariant quantum-mechanical spectra of hadrons (mesons and baryons) [2]. They are considered as two- and three-body systems of confined quarks interacting by different hyperfine interactions matching the experimentally measured energy levels [3]. While the spectra of quarks with only confinement – which has been often modelled by harmonic forces and is more realistically described by a linearly rising potential – are found to be regular, they exhibit chaotic behaviour upon the introduction of hyperfine interactions, motivated by quantum chromodynamics [4]. We consider hadron spectroscopy as a laboratory for studying quantum chaos, irrespective of the approach ranging from relativistic constituent-quark models [5] to lattice gauge theory [6].

References

- [1] T. Guhr, A. Müller-Groeling, and H.A. Weidenmüller: *Phys. Rep.* **299** (1998) 189
- [2] V. Pascalutsa: *Eur. Phys. J.* **16** (2003) 149
- [3] L.Y. Glozman, W. Plessas, K. Varga, and R.F. Wagenbrunn: *Phys. Rev. D* **58** (1998) 094030
- [4] H. Markum, W. Plessas, R. Pullirsch, B. Sengl, and R.F. Wagenbrunn: In Proceedings of the NATO Advanced Research Workshop on Nonlinear Dynamics and Fundamental Interactions, Tashkent, 2004; arXiv:hep-lat/0505011
- [5] W. Plessas: *Few-Body Syst. Suppl.* **15** (2003) 139
- [6] R. Pullirsch, K. Rabitsch, T. Wettig and H. Markum: *Phys. Lett. B* **427** (1998) 119

Semiclassical Paths to Many-Body Quantum Interference

Klaus Richter

Institut für Theoretische Physik, Universität Regensburg, Germany

Semiclassical methods in Quantum Chaos have been mainly successfully applied to single-particle quantum systems with corresponding low-dimensional chaotic dynamics. The generalization to interacting many-body systems remains to be one of the major challenges in this field. I will review first steps that have been made in my group [1-3] to devise semiclassical many-particle techniques. In particular, I will discuss a many-body representation of the Gutzwiller-Van Fleck propagator and show how this allows for investigating coherent backscattering and echo phenomena in Fock space, arising from many-body quantum interference.

References

- [1] Q. Hummel, J.-D. Urbina and K. Richter, The Weyl expansion for systems of independent identical particles, *J. Phys. A.* **47**, 01510 (2013).
- [2] T. Engl, J. Dujardin, A. Argüelles, P. Schlagheck, K. Richter and J.-D. Urbina, “Coherent Backscattering in Fock Space: a Signature of Quantum Many-Body Interference in Interacting Bosonic Systems”, *Phys. Rev. Lett.* **112**, 140403 (2014).
- [3] J.-D. Urbina, J. Kuipers, Q. Hummel, and K. Richter, “Multiparticle correlations in complex scattering and the mesoscopic Boson Sampling problem”, arXiv:1409.1558 [quant-ph]

Statistical properties of one-dimensional time-dependent Hamiltonian oscillators

Marko Robnik

CAMTP-Center for Applied Mathematics and Theoretical Physics, University of Maribor, Slovenia

Recently the interest in time-dependent dynamical systems has increased a lot. In this talk I shall present most recent results on time-dependent one-dimensional Hamiltonian oscillators. The time-dependence describes the interaction of an oscillator with its neighborhood. While the Liouville theorem still applies (the phase space volume is preserved), the energy of the system changes with time. We are interested in the statistical properties of the energy of an initial microcanonical ensemble with sharply defined initial energy, but uniform distribution of the initial conditions with respect to the canonical angle. We are in particular interested in the change of the action at the average energy, which is also adiabatic invariant, and is conserved in the ideal adiabatic limit, but otherwise changes with time. It will be shown that in the linear oscillator the value of the adiabatic invariant always increases, implying the increase of the Gibbs entropy in the mean (at the average energy). The energy is universally described by the arcsine distribution, independent of the driving law. In nonlinear oscillators things are different. For slow but not yet ideal adiabatic drivings the adiabatic invariant at the mean energy can decrease, just due to the nonlinearity and nonisochronicity, but nevertheless increases at faster drivings, including the limiting fastest possible driving, namely parametric kick (jump of the parameter). This is so-called PR property, following Papamikos and Robnik *J. Phys. A: Math. Theor.* **45** (2012) 015206, proven rigorously to be satisfied in a number of model potentials, such as homogeneous power law potential, and many others, giving evidence that the PR property is always satisfied in a parametric kick, except if we are too close to a separatrix or if the potential is not smooth enough. The local analysis is possible and the PR property is formulated in terms of a geometrical criterion for the underlying potential. We also study the periodic kicking and the strong (nonadiabatic) linear driving of the quartic oscillator. In the latter case we employ the nonlinear WKB method following Papamikos and Robnik *J. Phys. A: Math. Theor.* **44** (2012) 315102 and calculate the mean energy and the variance of the energy distribution, and also the adiabatic invariant which is asymptotically constant, but slightly higher than its initial value. The key references for the most recent work are Andreasas et al (2014), given below.

References

- Arnold V I 1980 *Mathematical Methods of Classical Mechanics* (New York: Springer-Verlag)
Lochak P and Meunier C 1988 *Multiphase Averaging for Classical Systems* (New York: Springer-Verlag)
Zaslavsky G M 2007 *The Physics of Chaos in Hamiltonian Systems* (London: Imperial College Press)
Ott E 1993 *Chaos in Dynamical Systems* (Cambridge University Press)
Chirikov B V 1979 *Phys. Rep.* **52** 263
Papamikos G and Robnik M 2011 *J. Phys. A: Math. Theor.* **44** 315102
Papamikos G, Sowden B C and Robnik M 2012 *Nonlinear Phenomena in Complex Systems (Minsk)* **15** 227
Papamikos G and Robnik M 2012 *J. Phys. A: Math. Theor.* **45** 015206
Robnik M and Romanovski V G 2006 *J. Phys. A: Math. Gen* **39** L35
Robnik M and Romanovski V G 2006 *Open Syst. & Infor. Dyn.* **13** 197
Robnik M, Romanovski V G and Stöckmann H.-J. 2006 *J. Phys. A: Math. Gen* L551
Kuzmin A V and Robnik M 2007 *Rep. on Math. Phys.* **60** 69
Robnik M V and Romanovski V G 2008 “Let’s Face Chaos through Nonlinear Dynamics”, Proceedings of the 7th International summer school/conference, Maribor, Slovenia, 2008, AIP Conf. Proc. No. 1076, Eds. M. Robnik and V.G. Romanovski (Melville, N.Y.: American Institute of Physics) 65
Robnik M and Romanovski V G 2000 *J. Phys. A: Math. Gen* **33** 5093
Andreasas D, Batistić B and Robnik M 2014 *Phys. Rev. E* **89** 062927
Andreasas D and Robnik M 2014 *J. Phys. A: Math. & Theor.* **46** 355102

- McLachlan R I 1995 *SIAM J.Sci.Comput.* **16** 151
- McLachlan R I and Quispel G R W 2002 *Acta Numerica* **11** 341
- Hairer E, Lubich C and Wanner G 2006 *Geometric Numerical Integration, Structure-Preserving Algorithms for Ordinary Differential Equations* (Berlin: Springer-Verlag)
- Leimkuhler B and Reich S 2004 *Simulating Hamiltonian Dynamics* (Cambridge: Cambridge University Press)
- Sanz-Serna J M and Calvo M P 1994 *Numerical Hamiltonian Problems* (London: Chapman & Hall)
- Shimada M and Yoshida H 1996 *Publ. Astron. Soc. Japan* **48** 147
- Yoshida H 1990 *Phys. Lett. A* **150** 262
- Yoshida H 1993 *Celestial Mechanics and Dynamical Astronomy* **56** 27
- Robnik M 2014 Time-dependent linear and nonlinear Hamilton oscillators *Selforganization in Complex Systems: The Past, Present and Future of Synergetics, Dedicated to Professor Hermann Haken on his 85th Anniversary (Proc. Int. Symp. Hanse Institute of Advanced Studies, Delmenhorst, 13-16 November 2012)* ed A Pelster and G Wunner (Berlin: Springer) at press
- Olver W F, Lozier W D, Boisvert F R and Clark W C 2010 *NIST Handbook of Mathematical Functions*, (Cambridge: Cambridge University Press)

Nonlinear Fourier transform and its inverse - a perturbational approach

Pavle Saksida

University of Ljubljana, Slovenia

The inverse scattering transform method for solving nonlinear integrable partial differential equations is a nonlinear analogue of the Fourier transform method for solving suitable initial-value problems for linear partial differential equations. Therefore, the scattering transform is often called the nonlinear Fourier transform (NLFT). When considering the inverse scattering transform method as a tool for finding solutions of initial problems, we find that it provides explicit analytic solutions only for certain very special initial conditions. The main difficulty resides in the fact that finding the inverse scattering transform (inverse NLFT) of a function is a difficult inverse problem, equivalent to a suitable Riemann-Hilbert problem. Therefore, it makes sense to resort to a perturbational approach. In the talk we shall describe a perturbational scheme by means of which we can explicitly compute the inverse NLFT of an arbitrary argument to any desired degree of accuracy. The convergence of this perturbational scheme is a consequence of certain analytical properties of the NLFT when defined on suitable function spaces. In the talk these properties will be proved.

Nested environments: Strong decoherence and coherence control

T. Gorin¹, H. Moreno², C. Gonzales² and **T. H. Seligman**^{2,3}

¹*Departamento de Física, Universidad de Guadalajara, Guadalajara, Mexico*

²*Instituto de Ciencias Físicas, Universidad Nacional Autónoma de México, Cuernavaca, Mexico*

³*Centro Internacional de Ciencias, Cuernavaca, Mexico*

For a central system weakly coupled to a near environment and uncoupled from a far environment The coupling between near and far environment can have surprising and possibly helpful effects on the conservation of coherence. We shall present analytic and numerical work in an RMT context as well as some numerics in spin networks. Finally we mention some exciting recent results of L. Campos Venuti and P. Zanardi on the use of such effects in coherence control.

Biological cell as a chronotaxic system

Aneta Stefanovska

Physics Department, Lancaster University, Lancaster, UK

The cell membrane is a lipid bilayer that divides the internal and external environment, thus making the cell the basic functioning unit of living systems. However, the cell is not independent – rather, a living cell continuously exchanges ions, oxygen and other nutrients with its environment known as the extracellular space. Charged ions are continuously crossing the cell membrane, actively and passively, thus resulting in a continuous change in the transmembrane potential. However, the membrane potential is traditionally considered to be in equilibrium. In this talk, based on experimental and theoretical work, we will discuss the origin of the fluctuations that are observed in the membrane potential and their time-dependent deterministic characteristics. In particular, we will argue that the cell can be described as a chronotaxic system (from *chronos* - time and *taxis* - order). This is a recently-introduced class of non-autonomous systems with time-dependent point attractors (driven steady states), able to resist external perturbations. The interplay between the production of ATP in mitochondria and due to glycolysis is modelled as a set of coupled chronotaxic systems. We will discuss the role of the oxygen supply and how the chronotaxic properties could change with cancer.

References

1. Y F Suprunenko, K Jenkins, G Lancaster, A Stefanovska, “Cell energy metabolism and its membrane potential”, in preparation.
2. J Jäckle, “The causal theory of the resting potential of cells”, *J Theor Biol* 249, 445-463 (2007).
3. D G Spiller, C D Wood, D A Rand, M R H White “Measurement of single-cell dynamics, *Nature* 465, 736-745 (2010).
4. A Goldbeter, C Gérard, D. Gonze, J-C Leloup, G Dupont, “Systems biology of cellular rhythms”, *FEBS Letters* 586, 2955–2965 (2012).
5. Y F Suprunenko, P Clemson, A Stefanovska, “Chronotaxic systems: A new class of self-sustained nonautonomous oscillators”, *Phys Rev Lett* 111, 024101 (2013).
6. P T Clemson, A Stefanovska, “Discerning non-autonomous dynamics”, *Phys Rep* (2014).
7. Y F Suprunenko, P T Clemson, A Stefanovska, “Chronotaxic systems with separable amplitude and phase dynamics”, *Phys Rev E* 89, 012922 (2014).
8. P T Clemson, Y F Suprunenko, T Stankovski, A Stefanovska, “The inverse approach to chronotaxic systems for single-variable time series”, *Phys Rev E* 89, 032904 (2014).

Microwave experiments in complex systems: From quantum chaos to monster waves

Hans-Juergen Stöckmann

Philipps-Universität Marburg, Germany

Most of the phenomena observed in wave propagation are universal and are observed in a large variety of different systems such as matter waves, but also, e.g., in electromagnetic waves and water waves. By means of the microwave measuring technique pioneered in Marburg, it thus becomes possible to study questions extending from the quantum mechanics of chaotic systems to the propagation of waves in the ocean. In this talk recent microwave results are presented, including the study of spectra and periodic orbits in microwave graphs and the study of branched flow in potential landscapes.

Classical and quantum chaos in the dynamics of molecules

Kazuo Takatsuka

Department of Basic Science, University of Tokyo, Tokyo, Japan

We present the following topics that we have studied and found, in which nonlinearity of dynamics manifests itself as characteristic phenomena of chaos in the dynamics of molecules.

- 1) Structural isomerization dynamics of atomic clusters: (a) loss of memory in the classical Hamilton chaos [1], (b) many-dimensional geometry behind the chaotic dynamics [2], (c) the concept of inter-basin mixing [3].
- 2) Bifurcation and merging of quantum wavepacket in consecutive nonadiabatic transitions [4]: (a) complexity arising from quantum entanglement between the dynamics of electrons and nuclei, (b) chaos in molecular double-slit experiments.
- 3) Chaos induced by quantum tunneling [5]: stationary states and their level statistics on a pair of coupled potential functions [6].
- 4) Induced photoemission from molecules of intra-molecular electron transfer: Chaotic fingerprints in photo-emission from a driven molecular oscillator (vibrational motion) of nuclear and electronic entanglement [7].
- 5) Phase-quantization of classical chaos [8]: mechanism of quantization of classical chaos.
- 6) Role of a newly found phase arising from the amplitude of a wavefunction in many-body quantum theory beyond semiclassics [9].

Among these, special emphases will be placed on the items (1), (4) and (6).

References:

- [1] K. Takatsuka and C. Seko, *J. Chem. Phys.* 105, 10356-10366 (1996).
- [2] K. Takatsuka, *Adv. Chem. Phys.* 130, 25-85 (2005).
- [3] K. Takatsuka and C. Seko, *J. Chem. Phys.* 110, 3263-3266 (1999)
- [4] Y. Arasaki, K. Takatsuka, K. Wang, and V. McKoy, *Phys. Rev. Lett.* 90, 248303 (2003).
- [5] H. Ushiyama and K. Takatsuka, *Phys. Rev. E* 53, 115-123 (1996).
- [6] H. Fujisaki and K. Takatsuka, *Phys. Rev. E* 63, 066221 (10 pages) (2001); H. Higuchi and K. Takatsuka, *Phys. Rev. E* 66, 035203(R) (4 pages) (2002).
- [7] Y. Arasaki, S. Scheit, K. Takatsuka, *J. Chem. Phys.*, 138, 161103 (4 pages) (2013).
- [8] S. Takahashi and K. Takatsuka, *J. Chem. Phys.* 127, 084112 (13 pages) (2007).
- [9] S. Takahashi and K. Takatsuka, *Phys. Rev. A* **89**, 012108 (2014); K. Takatsuka and S. Takahashi, *Phys. Rev. A* **89**, 012109 (2014).

Universal Quantum Graphs

Hans A. Weidenmueller

Heidelberg University

For completely connected simple graphs with incommensurate bond lengths and with unitary or orthogonal symmetry we prove the Bohigas-Giannoni-Schmit conjecture in its most general form. For graphs that are classically mixing (i.e., for which the spectrum of the classical Perron-Frobenius operator possesses a finite gap), we show that the generating functions for all (P, Q) correlation functions for both closed and open graphs coincide (in the limit of infinite graph size) with the corresponding expressions of random-matrix theory.

Quantum master equations: their power and limitations

Marko Znidaric

University of Ljubljana, Slovenia

Quantum master equations, frequently called Lindblad equations, have been used to describe physical systems, e.g., in quantum optics or NMR, long before general mathematical theory has been developed in '70 by Kossakowski, Sudarshan, Gorini, and Lindblad. In recent years Lindblad equations have become of interest also in statistical physics, condensed matter, and quantum information. On a practical side, Lindblad equations can be used as an efficient tool to study nonequilibrium physics of many-body systems. Theoretically they can be considered as a resource with which one can prepare quantum states, perform transformations, etc... While Lindblad equations can describe quite a large class of systems, they nevertheless have limitations and an interesting question one may ask is, which operations (or states) are possible to achieve with given resources? I will present some results in this respect as well as point out outstanding problems.

Music Concert



Nikolaj Sajko, cello
Miha Haas, piano

Program:

Ludwig van Beethoven: Sonata for piano and cello in A-major, Op. 69

- I. Allegro, ma non tanto
- II. Scherzo. Allegro molto
- III. Adagio cantabile - Allegro vivace

Lucijan Marija Škerjanc: Five Lyric Melodies for cello and piano

- I. Andante
- II. Lento
- III. Moderato cantabile
- IV. Lento
- V. Lento

Astor Piazzolla: Le Grand Tango

Nikolaj Sajko graduated in cello performance from the Academy of Music in Ljubljana in the class of Ciril Škerjanec and completed his Master of Science at the same institution, researching the articulation and phrasing of the orchestral playing. As a soloist he perfected his performance further at postgraduate studies at the Anton Bruckner Private University in Linz. During his studies he regularly took part in several international seminars with prominent professors such as I. Gavriš. E. Schoenfeld, A. Noras, C. Onczay and T. Kühne. Nikolaj Sajko was a member of the Gustav Mahler youth orchestra which operates under the leadership of Claudio Abbado. Besides that he was a member of the World Youth Orchestra. As a soloist Nikolaj performed with Maribor Philharmonic Orchestra, chamber orchestras Gaudeamus and Amadeus and with Maribor Youth Orchestra. He has performed in several solo recitals, chamber concerts and successful musical projects on the national as well as the international level. He was awarded the Klasinc diploma and Antonio Tarsia prize; other important achievements come from national and international competitions for young musicians in Gorizia, Lienz, Zagreb and Ljubljana. At present he is assistant solo cellist in the Symphony Orchestra of Slovene National Theatre in Maribor and a professor at the Conservatory for Music and Ballet in Maribor. In January this year Nikolaj released his first duo CD, along with pianist Miha Haas, with works for cello and piano by Edvard Grieg, Risto Savin, Bohuslav Martinů and Blaž Pucihar. Since 2012 Nikolaj is artistic leader of

chamber music series *Carpe artem*, which is co-organized by Musicians' Society Amadeus and Slovene National Theatre Maribor.

Miha Haas graduated in piano performance from the Academy of Music in Ljubljana in the class of Dubravka Tomšič-Srebotnjak and finished his post-graduation at The Royal Music Conservatory in Brussels in the class of Aleksandar Madžar. He took part at the international music seminars for piano and chamber music and worked with I. Lasko, J. Siirala, P. Paleczny, J. M. Huizing, M. Voskresenski, P. Kuijken and S. Picard. As a soloist he played with the Slovenian Philharmonic Orchestra, The RTV Slovenia Symphony Orchestra, The Symphonic Orchestra of the Academy of Music Ljubljana, The Symphonic Orchestra of Ljubljana Secondary School of Music, Celje String Orchestra, The Wind Orchestra of the Slovenian Army, The Wind Orchestra Ravne, and Festival Bled Orchestra. Moreover, his musical activities include also solo recitals, chamber music concerts especially in piano duos and in duos with the violin or cello. He participated in premier performances of 8 compositions by Slovenian composers and he made several recordings for the national radio and television. He was awarded the Prešeren prize of the Academy of Music Ljubljana and the Prešeren prize of the University of Ljubljana. Together with the pianist Tadej Horvat with whom he collaborates in a piano duo he won the second prize and the special prize for the best performance of the piece by E. Grieg at the Concours Grieg 2009 in Oslo. At present he works at the Academy of Music in Ljubljana.