
CAMTP

CENTER FOR APPLIED MATHEMATICS AND THEORETICAL PHYSICS
UNIVERZA V MARIBORU

13. Simpozij fizikov Univerze v Mariboru

Zbornik povzetkov

Hotel Piramida
Maribor, 11., 12. in 13. december 2014

Naslov: 13. Simpozij fizikov Univerze v Mariboru -
Zbornik povzetkov

Urednika: prof. dr. Marko Robnik
doc. dr. Anita Prapotnik Brdnik

Založnik/
Prireditelj simpozija: CAMTP - Center za uporabno
matematiko in teoretično fiziko,
Univerza v Mariboru

Kraj založbe: Maribor

Datum izida: December, 2014

Naklada: 50

Tisk: CopyCenter Oberčkal Bojan s.p.

CIP - Kataložni zapis o publikaciji
Univerzitetna knjižnica Maribor

53(082)(048.3)

SIMPOZIJ fizikov (13 ; 2014 ; Maribor)
Zbornik povzetkov / 13. simpozij fizikov Univerze v Mariboru, Maribor, 11., 12. in 13.
december 2014 ; [urednika Marko Robnik, Anita Prapotnik Brdnik]. - Maribor : CAMTP,
2014

ISBN 978-961-281-662-9
1. Robnik, Marko, 1954-

CAMTP

Organizacija simpozija: CAMTP - Center za uporabno matematiko in teoretično fiziko, Univerza v Mariboru

Organizacijski odbor:

Prof.Dr. Marko Robnik, CAMTP

Doc.Dr. Anita Prapotnik Brdnik, Katedra za aplikativno fiziko, Fakulteta za gradbeništvo

Urednika:

Prof.Dr. Marko Robnik, CAMTP

Doc.Dr. Anita Prapotnik Brdnik, Katedra za aplikativno fiziko, Fakulteta za gradbeništvo

Generalni sponzor simpozija:



E N E R G I J A

Generalni pokrovitelj simpozija:

EASA - European Academy of Sciences and Arts (Salzburg)



PREDGOVOR

Naši Simpoziji fizikov Univerze v Mariboru, ali na kratko kar Božični simpoziji, imajo že tradicijo, saj imamo letos že trinajstega po vrsti. Namen je strokovno druženje slovenskih fizikov, ob prisotnosti ter aktivni udeležbi nekaterih uglednih kolegov iz tujine kot častnih vabljenih gostov. Letos imamo šest uglednih vabljenih predavateljev iz tujine, iz odličnih raziskovalnih skupin, tako da s tem ostajajo naša srečanja nacionalna z mednarodno udeležbo. Srečanje je le ena od številnih dejavnosti CAMTP - Centra za uporabno matematiko in teoretično fiziko Univerze v Mariboru, ki sicer organizira kar sedem serij mednarodnih znanstvenih srečanj. Glej www.camtp.uni-mb.si

Radi bi poudarili, da je naše srečanje posvečeno vsej fiziki, teoretični in eksperimentalni, pa tudi matematični fiziki in uporabni matematiki in vsem drugim temam, za katere je fizika pomembna, ali pa so pomembne za fiziko.

Vsa predavanja so na ravni kolokvijev, se pravi razumljiva za splošnega fizika in zato še posebej primerna za študente, dodiplomske in poddiplomske. Takšnih splošnih srečanj na področju fizike v svetu pravzaprav skorajda ni več, čeprav so po našem prepričanju pomembna za širjenje intelektualnega obzorca vseh fizikov. Kolegi iz tujine, dosedanji udeleženci, potrjujejo to stališče in cenijo naš znanstveni program. Simpozij daje priložnost mladim raziskovalcem, da predstavijo svoje delo ter se o svojih rezultatih pogovorijo z izkušenimi znanstveniki. S to dejavnostjo prispevamo tudi k popularizaciji fizike v naši družbi na trajen način. Menimo, da je nujno poskrbeti za večjo popularizacijo naravoslovnih ved v naši družbi in fizika igra pri tem ključno vlogo. Vsem dodiplomskim študentom dovoljujemo brezplačno udeležbo na vseh predavanjih in s tem prispevamo k popularizaciji fizike ter k dodatnemu izobraževanju na tem področju.

Nenazadnje bi radi poudarili, da je naše druženje pomemben prispevek pri nadaljnjih uspešnih aktivnostih Fakultete za naravoslovje in matematiko Univerze v Mariboru ter Fakultete za matematiko in fiziko Univerze v Ljubljani in seveda institutov z oddelki za fiziko, kot sta Inštitut Jošef Stefan v Ljubljani ter CAMTP. V čast nam je, da bo med nami kot udeleženka tudi Dekanica FNM Prof. Dr. Nataša Vaupotič.

V čast nam je, da je generalni pokrovitelj našega srečanja Evropska akademija znanosti in umetnosti (European Academy of Sciences and Arts, Salzburg).

ORGANIZATORJA:

Prof.Dr. Marko Robnik, Direktor CAMTP
Doc.Dr. Anita Prapotnik Brdnik, FG UM

FOREWORD

Our Symposia of Physicists at the University of Maribor, or shortly Christmas Symposia, already have a tradition, as this year it is already the 13th one. The purpose is the scientific socializing of Slovenian physicists along with the participation of some distinguished colleagues from abroad as our honorary guests. This year we have six invited speakers from abroad, from some best research groups, so that our meetings remain national with international participation. The meeting is only one of the many activities of CAMTP - Center for Applied Mathematics and Theoretical Physics, which organizes seven series of international scientific meetings. See www.camtp.uni-mb.si

We would like to stress that our meeting is devoted to the entire physics, theoretical and experimental, and also applied mathematics and to all other topics, for which physics is important, or they are important for physics.

All lectures are on the level of colloquia, thus understandable for a general physicist, and therefore particularly well suited for students, the undergraduate and graduate students. Such general meetings in the field of physics practically no longer exist in the world, although to our opinion they are important for the widening of the intellectual horizon of all physicists. Our colleagues from abroad, the participants so far, confirm our view and appreciate our scientific programme. The meeting is also an opportunity for the young researchers to present their work and discuss it with the experienced scientists. With this activity we also contribute to the promotion and the popularization of physics in our society. We are convinced that it is quite urgent to care about the more intense popularization of natural sciences in our society, and physics plays a key role in this context. All undergraduate students can attend all the lectures of the conference free of charge. In this way we contribute to the popularization of physics and to the additional education in this field.

At the end we would like to stress that our gatherings are an important contribution to the activities of the Faculty of Natural Sciences and Mathematics of the University of Maribor and the Faculty of Mathematics and Physics of the University of Ljubljana, and of course also for the institutes with the department of physics, like Institute Jožef Stefan and CAMTP. It is our privilege that among the participants will be the Dean of FNM Prof. Dr. Nataša Vaupotič.

Our general patron is the European Academy of Sciences and Arts, Salzburg.

ORGANIZERS:

Prof.Dr. Marko Robnik, Direktor CAMTP
Prof.Dr. Anita Prapotnik Brdnik, FG UM

Seznam vseh udeležencev 13. Simpozija fizikov Univerze v Mariboru

Mr. Dimitrios Andrasas
CAMTP, University of Maribor
dimitraklos@hotmail.com

Prof.Dr. Denis Arčon
Institute Jožef Stefan, Ljubljana
denis.arcon@ijs.si

Mr. Iztok Bajc
FMF, University of Ljubljana
iztok.bajc@fmf.uni-lj.si

Mr. Benjamin Batistić
CAMTP, University of Maribor
benjamin.batistic@gmail.com

Prof.Dr. Tamas Biro
HAS Wigner Research Centre for Physics
Budapest, Hungary
Biro.Tamas@wigner.mta.hu

Prof.Dr. Janez Bonča
FMF, University of Ljubljana and IJS, Ljubljana
janez.bonca@fmf.uni-lj.si

Dr. Simon Čopar
FMF, University of Ljubljana
simon.copar@fmf.uni-lj.si

Ms. Maša Dukarić
CAMTP, University of Maribor
masa.dukaric@gmail.com

Prof.Dr. Rudolf Dvorak
Institute for Astronomy
University of Vienna, Austria
dvorak@astro.univie.ac.at

Prof.Dr. Svjetlana Fajfer
FMF, University of Ljubljana and IJS, Ljubljana
svjetlana.fajfer@ijs.si

Ms. Brigit Ferčec
CAMTP and FE, University of Maribor
brigit.fercec@gmail.com

Doc.Dr. Marko Gosak
FNM, University of Maribor
marko.gosak@uni-mb.si

Mr. Sašo Grozdanov
Instituut-Lorentz for Theoretical Physics
Leiden University, The Netherlands
grozdanov@lorentz.leidenuniv.nl

Mr. David Kraljič
University of Oxford, UK
David.Kraljic@physics.ox.ac.uk

Prof.Dr. Christian Lang
Institute for Theoretical Physics
University of Graz, Austria
christian.lang@uni-graz.at

Doc.Dr. Marjan Logar
FERI, University of Maribor
marjan.logar@um.si

Dr. Bing Sui Lu
FMF, University of Ljubljana
bing-sui.lu@fmf.uni-lj.si

Prof.Dr. Konstantinos Makris
Institute for Theoretical Physics
Technische Universität Wien, Austria
kgmakris78@gmail.com

Doc.Dr. Jernej Mravlje
Institute Jožef Stefan, Ljubljana
jernej.mravlje@ijs.si

Prof.Dr. Matjaž Perc
FNM and CAMTP, University of Maribor
matjaz.perc@uni-mb.si

Prof.Dr. Willibald Plessas
Institute for Theoretical Physics
University of Graz, Austria
willibald.plessas@uni-graz.at

Prof.Dr. Rudolf Podgornik
FMF, University of Ljubljana
rudolf.podgornik@fmf.uni-lj.si

Doc.Dr. Anita Prapotnik Brdnik
FG, University of Maribor
anita.prapotnik@um.si

Doc.Dr. Saša Prelovšek Komelj
FMF, University of Ljubljana and IJS, Ljubljana
sasa.prelovsek@ijs.si

Prof.Dr. Tomaž Prosen
Faculty of Mathematics and Physics, University of Ljubljana
tomaz.prosen@fmf.uni-lj.si

Prof.Dr. Marko Robnik
CAMTP, University of Maribor
robnik@uni-mb.si

Mr. Jakob Razdevšek
FNM, University of Maribor
jakob.razdevsek@gmail.com

Prof.Dr. Valery Romanovski
CAMTP, University of Maribor
valery.romanovsky@uni-mb.si

Prof.Dr. Douglas Shafer
Department of Mathematics
University of North Carolina
Charlotte, USA
dsshafer@uncc.edu

Mr. Vito Šimonka
FNM, University of Maribor
vito.simonka@gmail.com

Mr. Marko Šterk
FNM, University of Maribor
marko_sterk@hotmail.com

Prof.Dr. Nataša Vaupotič
FNM, University of Maribor
natasa.vaupotic@uni-mb.si

**Urnik 13. Simpozija fizikov
Univerze v Mariboru**

Četrtek, 11. december 2014	
Chair	Robnik
09:00-10:00	Podgornik
10:00-10:30	B. S. Lu
10:30-11:15	Mravlje
11:15-11:45	Coffee & Tea
11:45-12:30	Biro
12:30-13:15	Perc
13:15-15:00	Lunch
Chair	Prosen
15:00-15:45	Bonča
15:45-16:30	Fajfer
16:30-17:00	Coffee & Tea
17:00-17:45	Prapotnik
17:45-18:30	Batistić
19:00-20:00	Concert
20:00-23:00	Dinner

Petek, 12. december 2014	
Chair	Biro
09:00-10:00	Prosen
10:00-10:45	Dvorak
10:45-11:15	Andresas
11:15-11:45	Coffee & Tea
11:45-12:30	Robnik
12:30-13:15	Arčon
13:15-15:00	Lunch
Chair	Dvorak
15:00-15:45	Lang
15:45-16:30	Prelovšek
16:30-17:00	Coffee & Tea
17:00-17:45	Plessas
17:45-18:30	Grozdanov
18:30-18:45	Kraljič
19:00-20:00	Concert
20:00-23:00	Dinner

Sobota, 13. december 2014	
Chair	Arčon
09:00-09:45	Makris
09:45-10:30	Gosak
10:30-11:15	Čopar
11:15-11:45	Coffee & Tea
11:45-12:15	Vaupotič
12:15-13:00	Romanovski
13:00-13:15	Šimonka
13:15-15:00	Lunch
Chair	Plessas
15:00-15:45	Shafer
15:45-16:30	Ferčec
16:30-17:00	Coffee & Tea
17:00-17:30	Dukarić
17:30-17:45	Bajc
18:00-21:00	Dinner

Statistical properties of 1D time-dependent Hamilton systems

DIMITRIOS ANDRESAS, BENJAMIN BATISTIĆ and MARKO ROBNIK

CAMTP - *Center for Applied Mathematics and Theoretical Physics,*
University of Maribor, Maribor, Slovenia
dimitraklos@hotmail.com

I shall discuss studies of general nonlinear time-dependent 1D Hamiltonian systems. I shall present the general Papamikos-Robnik (PR) conjecture for parametrically kicked Hamilton systems, which says that for such systems the adiabatic invariant (the action) for an initial microcanonical ensemble at the mean final energy always increases under a parametric kick. I shall also present the ABR property, which is a local property, and many examples of the validity of this property which is almost always satisfied, but can be broken in not sufficiently smooth potentials or in cases where we are in the energy range close to a separatrix in the phase space. The general conjecture, using analytical and numerical computations, is shown to hold true for important systems like homogeneous power law potentials, pendulum, Kepler system, Morse potential, Pöschl-Teller I and II potentials, cosh potential, quadratic-linear potential, quadratic-quartic potential, while in three cases we demonstrate the absence of the ABR property: Linear oscillator enclosed in a box, sextic potential and quartic double well potential. In the second part of the talk I shall present the results of other kinds of time-dependent systems, assuming microcanonical ensemble of initial conditions, namely the cases of almost adiabatic (almost infinitely slow) variation, the case of unlimited linear driving of homogeneous power law potentials, where the nonlinear WKB-like method developed by Papamikos and Robnik (2012) can be applied, and analytic results on the statistical properties can be obtained. Finally, I shall discuss the cases of periodic driving.

References

- [1] G. Papamikos, B.C. Sowden and M. Robnik 2012 *Nonlinear Phenomena in Complex Systems (Minsk)* **15** 227
- [2] M. Robnik and V.G. Romanovski 2000 *J. Phys. A: Math. Gen* **33** 5093
- [3] M. Robnik and V.G. Romanovski 2006 *Open Syst. & Infor. Dyn.* **13** 197-222
- [4] G. Papamikos and M. Robnik 2011 *J. Phys. A: Math. Theor.* **44** 315102
- [5] G. Papamikos and M. Robnik 2012 *J. Phys. A: Math. Theor.* **45** 015206
- [6] D. Andrasas, B. Batistić and M. Robnik 2014 *Phys. Rev. E*, **89** 062927
- [7] D. Andrasas and M. Robnik 2014 *J. Phys. A: Math. Theor.*, **47** 355102

Superprevodnost in magnetizem v bližini prehoda kovina-izolator

DENIS ARČON

*Inštitut Jožef Stefan
F*

*Fakulteta za matematiko in fiziko, Univerza v Ljubljani
Jadranska 19, SI-1000 Ljubljana, Slovenia
denis.arcon@ijs.si*

Prehode kovina-izolator lahko v splošnem razdelimo v dve kategoriji: v prvi se izolatorsko stanje pojavi kot posledica neke strukturne spremembe, ki odpre energijsko režo med valenčnim in prevodnim pasom, v drugi pa je izvor prehoda povsem elektronski. To drugo kategorijo prehodov znova lahko razdelimo v dve podkategoriji: v prvo sodijo materiali, kjer je prehod posledica močnih elektronskih korelacij (t.i. Mottov prehod), v drugi pa je posledica nereda (t.i. Andersonov prehod).

V tem prispevku bom obravnaval obnašanje različnih elektronskih stanj v bližini takih prehodov. Najprej se bom posvetil Mottovem prehodu v fulerenskih soleh, kjer smo preko meritev z jedrsko magnetno resonanco pod visokimi pritiski spremeljali kako (antiferomagnetno) Mottovo-izolatorsko stanje zamenja superprevodnost s številnimi nenavadnimi lastnostmi [1]. V nadaljevanju pa bomo obravnavali vlogo nereda v Ti(O,N) nanožičkah, ki se neposredno odraža v transportnih lastnostih tako v normalni kot tudi v superprevodni fazi [2].

Reference

- [1] A. Potočnik et al., *Scientific Reports* **4** (2014) 4265.
- [2] D. Arčon et al., *neobjavljen*.

Superconductivity and magnetism in the vicinity of metal-to-insulator transition

DENIS ARČON

Institute Jožef Stefan
‐‐‐‐‐

*Faculty of mathematics and physics, University of Ljubljana
Jadranska 19, SI-1000 Ljubljana, Slovenia
denis.arcon@ijs.si*

Generally, metal-to-insulator transitions (MIT) can be divided into two categories. In the first category, the structural phase transition leads to a splitting of the electronic conduction band and hence to MIT. In the second category the transition is purely electronic in origin and can be again divided into two subcategories: one in which the transition is triggered by electronic correlations (i.e. Mott transition) and one in which it is triggered by disorder (i.e. Anderson transition).

In this contribution the behavior of competing electronic states in the vicinity of such transitions will be reviewed. In particular we will describe Mott transition in fulleride salts, where we have shown through comprehensive nuclear magnetic resonance measurements under high hydrostatic conditions that (antiferromagnetic) Mott insulating states is replaced by an unconventional superconductivity [1]. In the last part of the talk we will discuss the role of disorder in Ti(O,N) nanowires and how such disorder affects their transport properties in the normal and superconducting state [2].

References

- [1] A. Potočnik et al., *Scientific Reports* **4** (2014) 4265.
- [2] D. Arčon et al., *unpublished*.

Metoda končnih elementov za teoretsko fiziko

IZTOK BAJC

*FMF - Fakulteta za matematiko in fiziko
Univerza v Ljubljani, Jadranska 19, SI-1000 Ljubljana, Slovenija
iztok.bajc@fmf.uni-lj.si*

Predstavil bom nekaj izmed trenutnih zmogljivosti metode končnih elementov (FEM) za teoretsko fiziko.

S primerno vklopitvijo mrežne adaptivnosti v nelinearno minimizacijsko shemo lahko dosežemo v naprej nastavljeni napako. To smo uporabili pri minimizaciji Landau-de Gennes-ovega funkcionala proste energije in testirali na sistemih nematskih koloidov, a podobno je možno tudi za druge funkcionalne iz teoretske fizike, kot npr. Ginzburg-Landau za superprevodnost, ali Gross-Pitaevskij za Bose-Einsteinove kondenzate. Problem kako konkretno upoštevati brezdivergenčni pogoj za električna/magnetna polja v elektromagnetizmu ni bil zadovoljivo rešen kar nekaj let. Nédélec-ov tip končnega elementa je bil razvit nalašč za ta problem. Kratka koda, napisana v FreeFem++, točno izračuna lastne elektromagnetne načine v resonatorju napolnjenim z nehomogeno anizotropičnim materialom (nematski tekoči kristali).

Reference

- [1] I. Danaila and F. Hecht, A finite element method with mesh adaptivity for computing vortex states in fast-rotating Bose-Einstein condensates. *Jour. of Computat. Phys.* **229** (2010) 6946-6960.
- [2] Iz. Bajc, F. Hecht and S. Žumer, A mesh adaptivity scheme on the Landau-de Gennes functional minimization case in 3D, and its parametric efficiency (*to be submitted*).
- [3] I. Muševič, M. Škarabot, U. Tkalec, M. Ravnik and S. Žumer, Two-dimensional nematic colloidal crystals self-assembled by topological defects. *Science* (2006), **313**, 954.
- [4] G. E. Fasshauer, E. C. Gartland, Jr, J. W. Jerome, Newton iteration for partial differential equations and the approximation of the identity. *Numerical Algorithms* (2000), **25**, 181-195.

The finite element method for theoretical physics

IZTOK BAJC

*FMF - Faculty of Mathematics and Physics
University of Ljubljana, Jadranska 19, SI-1000 Ljubljana, Slovenia
iztok.bajc@fmf.uni-lj.si*

I will present some of the actual possibilities of the finite element method (FEM) for theoretical physics.

Mesh adaptivity, coupled to a nonlinear minimization scheme, can be used to reach an a priori fixed error threshold. This has been applied within the minimization of the Landau-de Gennes free energy functional for nematic liquid crystals, and tested on systems of nematic colloids, but similar functionals from theoretical physics, like the Ginzburg-Landau for superconductivity, or the Gross-Pitaevskij for Bose-Einstein condensates, apply as well.

How to impose the divergence-free condition for electric/magnetic fields in electromagnetism has not been satisfactorily solved/clarified for a couple of decades. The Nédélec type of finite element has been developed on porpose to solve this particular problem. A short code in FreeFem++ correctly computes EM eigen modes in a resonator filled with nonhomogeneously anisotropic material (nematic).

References

- [1] I. Danaila and F. Hecht, A finite element method with mesh adaptivity for computing vortex states in fast-rotating Bose-Einstein condensates. *Jour. of Computat. Phys.* **229** (2010) 6946-6960.
- [2] Iz. Bajc, F. Hecht and S. Žumer, A mesh adaptivity scheme on the Landau-de Gennes functional minimization case in 3D, and its parametric efficiency (*to be submitted*).
- [3] I. Muševič, M. Škarabot, U. Tkalec, M. Ravnik and S. Žumer, Two-dimensional nematic colloidal crystals self-assembled by topological defects. *Science* (2006), **313**, 954.
- [4] G. E. Fasshauer, E. C. Gartland, Jr, J. W. Jerome, Newton iteration for partial differential equations and the approximation of the identity. *Numerical Algorithms* (2000), **25**, 181-195.

Difuzija hitrosti adiabatskega biljardnega plina

BENJAMIN BATISTIĆ

CAMTP - Center for Applied Mathematics and Theoretical Physics

University of Maribor, Krekova 2, SI-2000 Maribor, Slovenia

benjamin.batistic@uni-mb.si • www.camtp.uni-mb.si

Obravnaval bom dinamiko porazdelitve hitrosti idealnega neinteragirajočega plina ujetega v časovno odvisnem oscilirajočem biljardu. Pokazal bom, da povprečna hitrost plina narašča eksponentno hitro v času, v kolikor je dinamika delca v statičnih biljardih, ki konstituirajo časovo odvisni biljard, mešanega tipa. To je precej drugače kot v plinu trdih kroglic, katerega porazdelitev hitrosti je opisana z Maxwell - Boltzmannovo porazdelitvijo in katerega povprečna hitrost je v adiabatski limiti omejena z adiabatsko invarianto. Pokazal bom, da Maxwell-Boltzmannova porazdelitev ne opiše porazdelitve hitrosti niti v primeru, ko je dinamika delca v biljardu popolnoma kaotična. Konstruiral bom Fokker-Planckovo enačbo za porazdelitev hitrosti in izpeljal njeno fundamentalno in asimptotsko rešitev, ter pokazal odlično ujemanje z numeričnimi rezultati.

Velocity diffusion in adiabatic billiard gas

BENJAMIN BATISTIĆ

*CAMTP - Center for Applied Mathematics and Theoretical Physics
University of Maribor, Krekova 2, SI-2000 Maribor, Slovenia
benjamin.batistic@gmail.com • www.camtp.uni-mb.si*

I will discuss the evolution of the distribution of velocities of the ideal noninteracting gas in a time-dependent oscillatory billiard. I will show that the average velocity of the gas increases exponentially fast in time if the particle dynamics in a corresponding static billiards of a time-dependent billiard is of the mixed type. This is very different from the gas of hard-balls whose velocity distribution is described by the Maxwell-Boltzmann distribution and whose average velocity is constrained by the adiabatic invariant in the adiabatic limit. I will show that the Maxwell-Boltzmann distribution does not apply even in a fully chaotic time-dependent billiards. I shall construct a Fokker-Planck equation for the velocity distribution in a fully chaotic time-dependent billiard and derive its fundamental and asymptotic solutions, and show that they are in perfect agreement with numerical results.

How to make your own entropy formula?

TAMÁS S. BIRÓ

*HAS Wigner Research Centre for Physics
Konkoly-Thege M ut 29-33, H-1221 Budapest, Hungary
Biro.Tamas@wigner.mta.hu • www.rmkki.kfki.hu/~tsbiro*

The good old exponential formula for the thermodynamical statistical weight factor at a given temperature, $\exp(-E/T)$, is valid only in the presence of huge (infinite) reservoirs in the background, keeping a sharp value for the subsystem's temperature. In high energy accelerator physics experiments typical reservoirs are small, they consist of a dozen or several hundreds to thousands of particles. The one particle energy spectra therefore are not exponential, even if treated statistically. We present statistical hadronization at a fixed total energy but varying particle number: the resulting picture of n -dimensional phase space volumina averaged over a distribution of n provides a cut power-law distribution of the one-particle energy. This generalizes the canonical exponential factor and includes it in the limit when n is Poisson distributed. Based on this experience we obtain the cut power-law statistical factor with general reservoir model equation of state, i.e. entropy – energy relation, $S(E)$, and identify the famous q -factor of non-extensive thermodynamics. Deviations from the Gibbs-exponential are due to finite reservoir size and induced fluctuations in the thermodynamical beta parameter, $\beta = S'(E)$. For Gaussian "normal" fluctuations the Gibbs-exponential is reconstructed. In the general case, however $q \neq 1$, and the entropy is not additive even for ideal systems. We propose to use instead a function of it, $K(S)$, which composes additively between subsystems. From this requirement a prescriptive differential equation emerges for $K(S)$, whose solution pinpoints a certain entropy – probability formula. This fundates the recipe for making your own entropy formula.

References

- [1] T. S. Biro, *Is There a Temperature? Conceptual Challenges at High Energy, Acceleration and Complexity*. Springer Verlag (2011) pp 310.
- [2] T. S. Biro and A. Jakovac, *Physical Review Letters* **94** (2005) 132302.
- [3] T. S. Biro and P. Van, *Physical Review E* **83** (2011) 061187.
- [4] T. S. Biro, G. G. Barnafoldi and P. Van, *EPJ A* **49** (2013) 110.

Relaksacijska dinamika več-delčnih sistemov

JANEZ BONČA

*Fakulteta za matematiko in fiziko, Univerza v Ljubljani, SI-1000
Ljubljana, Slovenija*

*Institut J. Stefan, SI-1000 Ljubljana, Slovenija
janez.bonca@ijs.si • www-f1.ijs.si*

Predstavil bom fundamentalno študijo ene in dveh vrzeli v t - J -Holsteinovem modelu pod vplivom zunanjega električnega polja. Ob upoštevanju kvantno mehanske narave problema sledimo časovnemu razvoju sistema začenši z osnovnim stanjem ko vključimo električno polje ob času nič ter vse do stacionarnega stanja. V primeru ene vrzeli opazimo adiabatski režim, kateremu sledi režim linearne $I - V$ karakteristike pri vmesnih električnih poljih [1,2,3]. Pri visokih poljih se sistem nahaja v režimu negativne diferencialne upornosti. Nadalje bom obravnaval tudi relaksacijo koreliranega sistema po vzbuditvi s kratkim električnim pulzom. Pokazal bom, da lahko foto vzbujeni nosilec naboja odda cca 1 eV energije spiskim eksitacijam v času reda velikosti nekaj fs [4]. V zaključku bom obravnaval relaksacijo dinamiko v primeru sklopitev s spiskimi ter mrežnimi prostostnimi stopnjami [5].

References

- [1] M. Mierzejewski, L. Vidmar, J. Bonča, and P. Prelovšek, *Phys. Rev. Lett.* **106** 196401, (2011).
- [2] L. Vidmar, J. Bonča, T. Tohyama, and S. Maekawa, *Phys. Rev. Lett.* **107** 246404, (2011).
- [3] J. Bonča, M. Mierzejewski, and L. Vidmar, *Phys. Rev. Lett.* **109** 156404, (2012).
- [4] D. Golež, J. Bonča, M. Mierzejewski, and L. Vidmar *Phys. Rev. B* **89** 165118 (2014).
- [5] J. Kogoj, Z. Lenarčič, D. Goleč, M. Mierzejewski, P. Prelovšek, and J. Bonča *Phys. Rev. B* **90**, 125104 (2014).

Relaxation dynamics of many-body systems

JANEZ BONČA

*Faculty of Mathematics and Physics, University of Ljubljana,
SI-1000 Ljubljana, Slovenia*

*J. Stefan Institute, SI-1000 Ljubljana, Slovenia
janez.bonca@ijs.si • www-f1.ijs.si*

I will present a fundamental study of one and two holes in the two dimensional t - J -Holstein model driven by the electric field. Taking fully into account quantum effects we follow the time-evolution of systems from their ground state as the electric field is switched on at $t = 0$, until they reach a steady state. In the single hole case adiabatic regime is observed followed by the positive differential resistivity at moderate fields where carrier mobility is determined [1,2,3]. At large field the system enters negative differential resistivity regime where current remains finite, proportional to $1/F$. I will also show that an ultrafast relaxation on the time-scale of 10 fs occurs due to inelastic scattering of a photo-excited carrier on spin excitations [4]. Finally, I will present relaxation dynamics of a carrier coupled to spin and lattice degrees of freedom [5].

References

- [1] M. Mierzejewski, L. Vidmar, J. Bonča, and P. Prelovšek, *Phys. Rev. Lett.* **106** 196401, (2011).
- [2] L. Vidmar, J. Bonča, T. Tohyama, and S. Maekawa, *Phys. Rev. Lett.* **107** 246404, (2011).
- [3] J. Bonča, M. Mierzejewski, and L. Vidmar, *Phys. Rev. Lett.* **109** 156404, (2012).
- [4] D. Golež, J. Bonča, M. Mierzejewski, and L. Vidmar *Phys. Rev. B* **89** 165118 (2014).
- [5] J. Kogoj, Z. Lenarčič, D. Goleč, M. Mierzejewski, P. Prelovšek, and J. Bonča *Phys. Rev. B* **90**, 125104 (2014).

Geometrijski gradniki topološke mehke snovi

SIMON ČOPAR

*Univerza v Ljubljani, Jadranska 19, SI-1000 Ljubljana, Slovenia
simon.copar@fmf.uni-lj.si*

Topološko zanimive teme, kot so stabilizacija defektov in solitonov, spletanje in vozlanje značilnih struktur, ter samourejevanje snovi, so uspešno raziskovalno področje v različnih panogah fizike. Med primere spadajo sinteza zavozlanih molekul [1], raziskave topologije v DNK in drugih polimerih [2], ter spletene strukture v ureditvenih poljih svetlobe [3], feromagnetov in različnih faz mehke snovi.

Vzorčni primer snovi, ki omogoča topološko stabilizirane strukture, so nematski tekoči kristali. Zaradi obstoja točkastih in linijskih defektov, močnega odziva na optično in električno polje, ter enostavnosti opazovanja pod mikroskopom, so idealni za demonstracijo teorije vozlov in interakcij med defekti v eksperimentalni postavitvi. Predstavl bom teoretične zakone spletanja in vozlanja nematskih defektov, ki uvedejo topološke invariante in pojasnijo strukture, videne v eksperimentih [4]. Prikazal bom obnašanje in nadzor spletanja defektov na 2D in 3D mrežah koloidnih delcev v tekočekristalnem mediju [5,6], ter v kapljicah kiralnega nematica [7]. Primeri se razlikujejo po geometriji prostora, ki ga zapoljuje tekoči kristal, in po simetrijskih lastnostih samega tekočega kristala. Predstavl bom temeljne geometrijske gradnike, ki sestavljajo splete defektov v posameznih primerih.

Reference

- [1] J. F. Ayme, J. E. Beves, D. A. Leigh, R. T. McBurney, K. Rissanen, D. Schultz, *J Am. Chem. Soc.* **134**, 9488-9497 (2012).
- [2] C. Micheletti, E. Orlandini, *ACS Macro Letters* **3**, 876-880 (2014).
- [3] M. R. Dennis, R. P. King, B. Jack, K. O. Holleran, M. J. Padgett, *Nature Physics* **6**, 118 - 121 (2010).
- [4] S. Čopar, *Phys. Rep.* **538**, 1-37 (2014).
- [5] U. Tkalec, M. Ravnik, S. Čopar, S. Žumer, I. Muševič, *Science* **333**, 62 (2011).
- [6] S. Čopar, N. A. Clark, M. Ravnik, S. Žumer, *Soft Matter* **9**, 8203 (2013).
- [7] D. Seč, S. Čopar, S. Žumer, *Nature Commun.* **5**, 3057 (2014).

Geometric building blocks in topological soft matter

SIMON ČOPAR

*Univerza v Ljubljani, Jadranska 19, SI-1000 Ljubljana, Slovenia
simon.copar@fmf.uni-lj.si*

Topological topics, such as stabilization of solitons and defects, knotting and linking of representative structures and self-assembly are targets of active development in different fields. Among them are targeted synthesis of knotted molecules [1], research of topological states in DNK and other polymers [2], and textures in ordering fields in light [3], ferromagnets and various soft materials.

Nematic liquid crystals are a typical medium that supports topologically stabilized structures. The existence of point and line defects, strong response to optical and electric fields, and easy observation under the microscope, make them ideal for demonstration of knot theory and defect interaction in experimental environment. I present theoretical laws of linking and knotting of nematic defects, introduce topological invariants and explain structures seen in experiments [4]. I demonstrate the behaviour of knotted defects in 2D and 3D colloidal grids in LC medium [5,6] and chiral LC droplets [7]. The cases differ in geometry of the space occupied by the liquid crystal and in symmetry properties of the liquid crystalline structure itself. I review the basic geometric building blocks of defect networks in each of the cases.

References

- [1] J. F. Ayme, J. E. Beves, D. A. Leigh, R. T. McBurney, K. Rissanen, D. Schultz, *J Am. Chem. Soc.* **134**, 9488-9497 (2012).
- [2] C. Micheletti, E. Orlandini, *ACS Macro Letters* **3**, 876-880 (2014).
- [3] M. R. Dennis, R. P. King, B. Jack, K. O. Holleran, M. J. Padgett, *Nature Physics* **6**, 118 - 121 (2010).
- [4] S. Čopar, *Phys. Rep.* **538**, 1-37 (2014).
- [5] U. Tkalec, M. Ravnik, S. Čopar, S. Žumer, I. Muševič, *Science* **333**, 62 (2011).
- [6] S. Čopar, N. A. Clark, M. Ravnik, S. Žumer, *Soft Matter* **9**, 8203 (2013).
- [7] D. Seč, S. Čopar, S. Žumer, *Nature Commun.* **5**, 3057 (2014).

Lokalna integrabilnost in linearizabilnost (1 : -1 : -1) resonantnega kvadratičnega sistema

MAŠA DUKARIĆ, VALERY G. ROMANOVSKI

CAMTP - Center za uporabno matematiko in teoretično fiziko
Univerza v Mariboru, Krekova 2, SI-2000 Maribor, Slovenija
masa.dukaric@gmail.si, valery.romanovsky@uni-mb.si

REGILENE OLIVEIRA

Univerza São Paulo, Departamento de Matemática, ICMC-USP
Caixa Postal 668, 13560-000 São Carlos, Brazilija
regilene@icmc.usp.br

Raziskovali smo kompletno integrabilnost in linearizabilnost tri-dimenzionalnega sistema oblike

$$\begin{aligned}\dot{x} &= x + a_{12}xy + a_{13}xz + a_{23}yz \\ \dot{y} &= -y + b_{12}xy + b_{13}xz + b_{23}yz \\ \dot{z} &= -z + c_{12}xy + c_{13}xz + c_{23}yz.\end{aligned}$$

Pridobljeni so bili potrebni in zadostni pogoji obstoja dveh neodvisnih analitičnih integralov. Potrebni pogoji so bili izračunani z uporabo algoritmov računske komutativne algebre. Za dokaz integrabilnosti in linearizabilnosti je bila uporabljena Darboux-jeva metoda ter teorija normalnih form. Nekateri Darboux-jevi faktorji potrebni za linearizabilnost sistema so bili pridobljeni iz prvih integralov sistema. Obravnavali smo tudi problem obstoja enega analitičnega prvega integrala.

Reference

- [1] W. Aziz, C. Christopher: *Local integrability and linearizability of three-dimensional Lotka-Volterra systems*, Appl. Math. Comput. **21** (2012), no. 8, 4067–4081.
- [2] W. Aziz: *Integrability and Linearizability of Three dimensional vector fields*, Qual. Theory of Dyn. Syst **13** (2014), 197–213.
- [3] Z.Hu, M. Han, V.G. Romanovski: *Local integrability of a family of three-dimensional quadratic systems*, Physica D: Nonlinear Phenomena **265** (2013), 78,1rs-86.

Local integrability and linearizability of a (1 : -1 : -1) resonant quadratic system

MAŠA DUKARIĆ, VALERY G. ROMANOVSKI

CAMTP- Center for Applied Mathematics and Theoretical Physics
University of Maribor, Krekova 2, SI-2000 Maribor, Slovenia
masa.dukaric@gmail.si, valery.romanovsky@uni-mb.si

REGILENE OLIVEIRA

University of São Paulo, Departamento de Matemática, ICMC-USP
Caixa Postal 668, 13560-000 São Carlos, Brazil
regilene@icmc.usp.br

We study the complete integrability and linearizability of a three dimensional system of the form

$$\begin{aligned}\dot{x} &= x + a_{12}xy + a_{13}xz + a_{23}yz \\ \dot{y} &= -y + b_{12}xy + b_{13}xz + b_{23}yz \\ \dot{z} &= -z + c_{12}xy + c_{13}xz + c_{23}yz.\end{aligned}$$

Necessary and sufficient conditions for existence of two functionally independent analytic integrals in the family are obtained. The necessary conditions are derived using algorithms of computational commutative algebra. To prove integrability and linearizability we used the method of Darboux and the normal form theory. Some Darboux factors used for linearizability are obtained from first integrals of systems. The problem of existence of only one analytic first integral is discussed as well.

References

- [1] W. Aziz, C. Christopher: *Local integrability and linearizability of three-dimensional Lotka-Volterra systems*, Appl. Math. Comput. **21** (2012), no. 8, 4067–4081.
- [2] W. Aziz: *Integrability and Linearizability of Three dimensional vector fields*, Qual. Theory of Dyn. Syst **13** (2014), 197–213.
- [3] Z.Hu, M. Han, V.G. Romanovski: *Local integrability of a family of three-dimensional quadratic systems*, Physica D: Nonlinear Phenomena **265** (2013), 78, 1rs-86.

Collisions of planetesimals and the formation of planets

Rudolf Dvorak, Thomas I. Maindl and Christoph Burger

University of Vienna, Institute of Astronomy

Since the last 20 years many planets (approx. 1850 up to now) around other stars were detected either from the ground or by space missions. More than 500 planetary systems with more than one discovered planet are known; for planets being habitable many conditions need to be fulfilled: e.g. the spectral type and age of the hosting star, the location in our galaxy and also the orbit of a planet and its physical nature. Different estimates lead to quite different results especially for planet which host in fact life. In this respect we show the results of collision computations with the aid of Smooth Particle Hydrodynamic (SPH) codes and discuss also in short the role of the tides acting on these bodies.

Reference

- [1] Dvorak R, Eggel S, Süli A, Sandor Z, Galiazzo M and Pilat-Lohinger, E 2012, *AIP Conference Proceedings* **1468** 137.
- [2] Maindl T I, Schäfer C, Speith R, Süli, A and Forgács-Dajka, E 2013, *Astronomische Nachrichten* **334** 996.
- [3] Maindl T I, Dvorak R, Speith R and Schäfer C 2013, *2014arXiv1401.0045M*, 12.
- [4] Maindl T I and Dvorak R 2014, *IAU Symposium* **299**, 370.
- [5] Schäfer C, Speith R, Hipp, M and Kley W 2004, *Astronomy and Astrophysics* **418**, 325.

Leptokvarki v fiziki okusov in hadronskih trkalnikov

SVJETLANA FAJFER

*Oddelek za fiziko FMF, Univerza v Ljubljani, Jadranska 19, SI-1000
Ljubljana, Slovenija*

*Institut J. Stefan, Jamova 39, SI-1000 Ljubljana, Slovenija
svjetlana.fajfer@ijs.si*

Leptoquarki interagirajo s kvarki in leptoni in so običajno prisotni v teorijah poenotenja osnovnih interakcij. V teh teorijah, praviloma, imajo mase blizu mase v skale GUT. Vendar, pa v nekaterih teorijah poenotenja imajo lahko skalarni leptokvarki mase reda velikosti 1 TeV. Takšna stanja so lahko prisotna v procesih na nizkih energijah. Obenem, jih lahko opazijo na hadronskih trkalnikih. Naš studij je posvečen raziskavi lahkih leptoquarkov, kateri ne morejo destabilizirati proton. Takšni ”nenevarni” leptokvarki lahko imajo relativno velike sklopitve s fermioni standardnega modela. Obravnavamo vpliv teh sklopitev na mehanizme produkcije leptokvarkov na LHC in ugotavljamo, da je za produkcijo leptokvarkov v teh primerih zelo pomemben prispevek t-kanala.

Reference

- [1] I. Doršner, S. Fajfer and A. Greljo, *JHEP* **1410** (2014) 154.
- [2] I. Doršner, S. Fajfer, N. Košnik and I. Nišandžić, *JHEP* **1311** (2013) 084.
- [3] I. Doršner, S. Fajfer, A. Greljo and J. F. Kamenik, *JHEP* **1211** (2012) 130.
- [4] I. DoršnerDorsner, S. Fajfer and N. Košnik, *Phys. Rev. D* **86** (2012) 015013.
- [5] I. DoršnerDorsner, S. Fajfer, J. F. Kamenik and N. Košnik, *Phys. Rev. D* **82** (2010) 094015.
- [6] I. DoršnerDorsner, S. Fajfer, J. F. Kamenik and N. Košnik, *Phys. Rev. D* **81** (2010) 055009.
- [7] I. Doršner, S. Fajfer, J. F. Kamenik and N. Košnik, *Phys. Lett. B* **682** (2009) 67.
- [8] S. Fajfer and N. Košnik, *Phys. Rev. D* **79** (2009) 017502.

Scalar Leptoquarks in Flavor and Collider Physics

SVJETLANA FAJFER

*Physics Department, University of Ljubljana, Jadranska 19, SI-1000
Ljubljana, Slovenia*

*Jozef Stefan Institute , Jamova 39, SI-1000 Ljubljana, Slovenia
svjetlana.fajfer@ijs.si*

Leptoquarks interact with quarks and leptons and they are usually present in theories which unify fundamental interactions. In these theories they naturally have a mass close to the GUT scale. However, in some of unification approaches scalar leptoquarks might have masses of the order 1 TeV. Such are states might modify processes at the low-energies. At the same time they can be seen at Large hadronic collider directly. These light leptoquark states should not mediate proton decay at the tree level. Our analysis is devoted to the light leptoquarks which do not destabilise proton. These "non-dangerous" scalar leptoquarks might have sizeable couplings to the Standard Model fermions. We discuss leptoquark production mechanisms at LHC and demonstrate the importance of inclusion of a t-channel pair production.

References

- [1] I. Doršner, S. Fajfer and A. Greljo, *JHEP* **1410** (2014) 154.
- [2] I. Doršner, S. Fajfer, N. Košnik and I. Nišandžić, *JHEP* **1311** (2013) 084.
- [3] I. Doršner, S. Fajfer, A. Greljo and J. F. Kamenik, *JHEP* **1211** (2012) 130.
- [4] I. DoršnerDorsner, S. Fajfer and N. Košnik, *Phys. Rev. D* **86** (2012) 015013.
- [5] I. DoršnerDorsner, S. Fajfer, J. F. Kamenik and N. Košnik, *Phys. Rev. D* **82** (2010) 094015.
- [6] I. DoršnerDorsner, S. Fajfer, J. F. Kamenik and N. Košnik, *Phys. Rev. D* **81** (2010) 055009.
- [7] I. Doršner, S. Fajfer, J. F. Kamenik and N. Košnik, *Phys. Lett. B* **682** (2009) 67.
- [8] S. Fajfer and N. Košnik, *Phys. Rev. D* **79** (2009) 017502.

Hopfove bifurkacije v sistemih NDE

BRIGITA FERČEC

*CAMTP - Center za uporabno matematiko in teoretično fiziko,
Univerza v Mariboru, Maribor, Slovenija
FE - Fakulteta za energetiko,
Univerza v Mariboru, Krško, Slovenija*

Najprej bom razložila Hopfovo (ali Andronov-Hopfovo) bifurkacijo za ravninske sisteme. Opisala bom limitne cikle na centralni mnogoterosti in degenerirane Hopfove (Bautinove) bifurkacije. Nato bom predstavila enega izmed najbolj znanih problemov kvalitativne teorije navadnih diferencialnih enačb - Hilbertov 16. problem o številu limitnih ciklov dvo-dimenzionalnih sistemov

$$\dot{x} = P_n(x, y), \quad \dot{y} = Q_n(x, y)$$

(n je maksimalna stopnja polinomov na desni strani sistema). Bistven del problema je problem ocenitve maksimalnega števila limitnih ciklov, ki bifurcirajo iz singularne točke tipa center ali fokus pod vplivom majhnih motenj koeficientov sistema, t.j. problem cikličnosti. Ključna značilnost našega pristopa je, da je v primeru elementarne singularne točke problem cikličnosti reducirana na algebraični problem iskanja baze določenega polinomskega idealja. Ta pristop nato uporabimo za problem cikličnosti poddržine kubičnih sistemov.

Reference

- [1] Ferčec B. and Mahdi A. 2013 *Math. Comput. Simul.* **87** 55–67.
- [2] Kuznetsov Yu.A. 2004 *Elements of Applied Bifurcation Theory* Springer - Verlag, New York.
- [3] Romanovski V. G. and Shafer D. S. 2009 *The center and cyclicity problems: A computational algebra approach* Birkhäuser, Boston.

Hopf bifurcations in systems of ODE's

BRIGITA FERČEC

*CAMTP - Center for Applied Mathematics and Theoretical Physics,
University of Maribor, Maribor, Slovenia
FE - Faculty of Energy Technology,
University of Maribor, Krško, Slovenia*

First, I will discuss a Hopf (or Andronov-Hopf) bifurcation for planar systems. I will describe also limit cycles on the center manifold and degenerate Hopf (Bautin) bifurcations. Then, I will present one of the most famous problems in qualitative theory of ordinary differential equations - Hilbert's sixteenth problem on the number of limit cycles of two dimensional polynomial systems

$$\dot{x} = P_n(x, y), \quad \dot{y} = Q_n(x, y)$$

(n is the maximum degree of the polynomials on the right-hand side of the system). An essential part of the problem is the problem of estimating of the maximum number of limit cycles which can bifurcate from a singular point of center or focus type under small perturbations of coefficients of the system, the so-called cyclicity problem. The key feature of our approach is that in the case of an elementary singular point the problem of cyclicity is reduced to the algebraic problem of searching for a basis of a certain polynomial ideal. We apply this approach for solving the cyclicity problem for a subfamily of cubic systems.

References

- [1] Ferčec B. and Mahdi A. 2013 *Math. Comput. Simul.* **87** 55–67.
- [2] Kuznetsov Yu.A. 2004 *Elements of Applied Bifurcation Theory* Springer - Verlag, New York.
- [3] Romanovski V. G. and Shafer D. S. 2009 *The center and cyclicity problems: A computational algebra approach* Birkhäuser, Boston.

Uporaba pristopov s področja kompleksnih sistemov za preučevanje funkcionalne in strukturne organiziranosti sincicija celic beta

Marko Gosak^{1,2}, Andraž Stožer³, Rene Markovič¹, Jurij Dolenšek³, Thanos Manos⁴, Marko Marhl^{1,2} in Marjan Slak Rupnik^{3,5}

¹*Fakulteta za naravoslovje in matematiko, Univerza v Mariboru,
Slovenija*

²*Pedagoka fakulteta, Univerza v Mariboru, Slovenija*

³*Inštitut za fiziologijo, Medicinska fakulteta, Univerza v Mariboru,
Slovenija*

⁴*Raziskovalni center Jülich, Nemčija*

⁵*Medicinska univerza na Dunaju, Avstrija*

Metodologija s področja kompleksnih sistemov se dandanes vse pogosteje uporablja za preučevanje in razumevanje organizacijskih ter funkcionalnih principov bioloških sistemov, in to na najrazličnejših skalah [1]. V naši študiji se osredotočimo na kolektivno dinamiko celic beta v Langerhansovih otočkih trebušne slinavke, ki vodi do izločanja inzulina. Na osnovi korelacij med izmerjenimi časovnimi vrstami znotrajceličnih kalcijevih signalov, pridobljenih s konfokalnim mikroskopom, smo izgradili funkcionalno mrežo celic beta, s katero okarakteriziramo medcelično komuniciranje [2]. Ekstrahirane vzorce medcelične povezanosti analiziramo z uveljavljenimi orodji s področja teorije kompleksnih mrež. Z izračunom različnih mrežnih mer si ne pridobimo le informacij o naravi funkcionalnih medceličnih povezav, temveč lahko tudi sledimo časovnemu razvoju strukture mreže [3]. Za pridobitev še bolj poglobljenega vpogleda v delovanje Langerhansovih otočkov se poslužimo tudi matematičnega modeliranja. Izdelali smo podrobni večelični matematični model sincicija celic beta, kjer posamezne celice predstavljajo vozle, povezave pa ponazarjajo medcelično komunikacijo. V model smo vključili tudi dodatne eksperimentalno ugotovljene fiziološke detajle. Naši rezultati pokažejo, da je kombinacija naprednih mikroskopskih tehnik, sodobnih analitičnih orodji in matematičnega modeliranja zelo udarno orodje, s katerim si lahko pridobimo poglobljen vpogled v delovanje Langerhansovih otočkov. Prav tako nam tovrstna metodologija nudi dobro izhodišče za preučevanje modifikacij povezanih s patofiziološkimi stanji, kot je na primer sladkorna bolezen.

Reference

- [1] A.-L. Barabási, *Nature Phys.* **8** (2012) 14.
- [2] A. Stožer, M. Gosak, J. Dolenšek, M. Perc, M. Marhl, M. S. Rupnik, D. Korošak, *PLoS Comput Biol* **9** (2013) e1002923.
- [3] R. Markovič, A. Stožer, M. Gosak, J. Dolenšek, M. Marhl, M. S. Rupnik, poslano v *Scientific Reports*.

Application of complex systems approaches for studying the functional and structural organization of the pancreatic beta cell syncytium

Marko Gosak^{1,2}, Andraž Stožer³, Rene Markovič¹, Jurij Dolenšek³,
Thanos Manos⁴, Marko Marhl^{1,2} and Marjan Slak Rupnik^{3,5}

¹*Faculty of Natural Sciences and Mathematics, University of
Maribor, Slovenia*

²*Faculty of Education, University of Maribor, Slovenia*

³*Faculty of Medicine, University of Maribor, Slovenia*

⁴*Research Center Jülich, Germany*

⁵*Medical University of Vienna, Austria*

Methodology developed in the field of complex systems science is nowadays frequently used for the analysis of the structure and function of biological systems at different scales [1]. In our study we focus on the collective pancreatic beta cell activity in the islets of Langerhans, which triggers and regulates insulin secretion. For the characterization of the intercellular communication we build up functional networks on the basis of correlations between calcium dynamics of individual cells, which were recorded by means of confocal laser-scanning microscopy [2]. The extracted pattern of pairwise interactions between network cells, i.e. functional connections, is then scrutinized with conventional tools for network analysis. By calculating the network metrics of interest, we can not only characterize the nature of functional connectivity patterns, but can also track and analyze the temporal evolution of tissue networks [3]. In order to complement our understanding of the mechanisms that govern the functioning of islets, we make use of mathematical modelling approaches. We propose a detailed multicellular model of interconnected beta cells, which upgrades the existing modeling endeavors by incorporating additional physiological specifics. Our results demonstrate that a combination of advanced high spatial and temporal resolution confocal imaging, novel analytical tools and computational modeling is a powerful tool that enables not only a comprehensive understanding of functional mechanisms and organization of islets of Langerhans but also provides a firm framework for the examination of modifications associated with pathophysiological conditions, such as diabetes mellitus.

References

- [1] A.-L. Barabási, *Nature Phys.* **8** (2012) 14.
- [2] A. Stožer, M. Gosak, J. Dolenšek, M. Perc, M. Marhl, M. S. Rupnik, D. Korošak, *PLoS Comput Biol* **9** (2013) e1002923.
- [3] R. Markovič, A. Stožer, M. Gosak, J. Dolenšek, M. Marhl, M. S. Rupnik, submitted to *Scientific Reports*.

Second-order hydrodynamics and universality from the gauge/string duality

SASO GROZDANOV

*Instituut-Lorentz for Theoretical Physics, Leiden University,
Niels Bohrweg 2, Leiden 2333 CA, The Netherlands
grozdanov@lorentz.leidenuniv.nl*

The gauge/string duality has provided us with a tool for analysing certain classes of strongly coupled field theories. In its simplest limit of a large number of colours and a large 't Hooft coupling, it enables us to compute field theory correlation functions at all energy scales, directly from classical gravity. In the hydrodynamical limit, the duality gives us both the structure of the hydrodynamical gradient expansion at all orders as well the transport coefficients of the fluid. We can therefore use it to study higher-order corrections to the one-derivative Navier-Stokes equations. After presenting the gauge/string duality, I will discuss second-order hydrodynamics and a universal result relating three of its transport coefficients at strong coupling. I will focus on higher-derivative gravity corrections to the universal relation, particularly in a non-perturbative limit in which viscosity vanishes and second-order hydrodynamics becomes dominant. Finally, the phenomenology of nearly dissipationless fluids will be discussed, using the holographic Gauss-Bonnet fluid as an example.

References

- [1] G. Policastro, D. T. Son and A. O. Starinets, “From AdS / CFT correspondence to hydrodynamics,” JHEP **0209** (2002) 043 [hep-th/0205052].
- [2] M. Haack and A. Yarom, “Universality of second order transport coefficients from the gauge-string duality,” Nucl. Phys. B **813** (2009) 140 [arXiv:0811.1794 [hep-th]].
- [3] S. Grozdanov and A. O. Starinets, “Gauss-Bonnet gravity and dissipationless fluids,” to appear.
- [4] S. Grozdanov and A. O. Starinets, “On the universal identity in second-order hydrodynamics,” to appear.
- [5] S. Grozdanov and A. O. Starinets, “The limit of zero viscosity in a holographic Gauss-Bonnet fluid,” Special issue of Theor. Math. Phys. in honor of A.A. Slavnov’s 75th birthday.

Jata galaksij ”Bullet” in ekstremni objekti v Hubblovem volumnu

DAVID KRALJIĆ

Rudolf Peierls Centre for Theoretical Physics

University of Oxford, 1 Keble Road, Oxford, UK

david.kraljic@physics.ox.ac.uk • www.physics.ox.ac.uk

Velike jate in nadjate galaksij so največji gravitacijsko vezani objekti v našem vesolju. Pričakujemo, da se zaradi širjenja vesolja premikajo stran drug od drugega. V zadnjih časih so bile odkrite številne jate, ki so v procesu trčenja - nasprotno od naivnega pričakovanja. Med njimi je najbolj raziskovana in slavna ”Bullet” jata galaksij. Torej, preveriti moramo ali je naš Λ CDM model vesolja zmožen generirati trke velikih jat galaksij. Ker so trki nelinearni, se predikcije Λ CDM modela testirajo v kontekstu kozmoloških simulacij N-teles. V preteklosti se je pojavil konsenz, da so trki kot je ”Bullet” jata malo verjetni in jih ne pričakujemo v našem vesolju. Opis in preučevanje ekstremnih dogodkov v repih distribucij zahteva nestandardna statistična orodja (npr. Statistika ekstremnih vrednosti) in precizno poznavanje numeričnih metod v simulacijah. Nove metode, skupaj z eno od največjih kozmoloških simulacij N-teles, nas pripeljejo do sklepa, da so trki jat galaksij bolj pogosti, kot smo pričakovali v preteklosti.

The Bullet cluster and extreme objects in the Hubble volume

DAVID KRALJIĆ

*Rudolf Peierls Centre for Theoretical Physics
University of Oxford, 1 Keble Road, Oxford, UK
david.kraljic@physics.ox.ac.uk • www.physics.ox.ac.uk*

Large galaxy clusters are the biggest gravitationally bound objects in our universe. They are expected to follow the Hubble flow - move away from each other due to the expansion of the universe. However, a significant number of colliding clusters has been found, with the Bullet cluster being the best studied and most famous. Therefore, our naive expectation and the concordance Λ CDM model have to be reconciled with the observation. Because of their highly non-linear nature, cluster collisions have been studied in the context of cosmological N-Body simulations and were found to be highly unlikely. Studying the extreme events in the tails of distributions requires non-standard statistical tools (e.g. Extreme Value Statistics) as well as a precise understanding of the numerical methods used in the simulations. Using improved methodology and one of the largest simulations of our universe we show that the collisions of large clusters do occur and are not as rare as previously thought.

Lattice hadron scattering: An introduction

CHRISTIAN B. LANG

*Institute of Physics
University of Graz, A-8010 Graz, Austria
christian.lang@uni-graz.at*

Recent years have seen considerable progress in ab initio QCD calculations of hadron scattering threshold parameters and scattering phase shifts in the (elastic) resonance region. The lattice approach is becoming powerful enough to even predict states in the heavy quark sector. I will discuss the methods and limitations in an introductory way, In a subsequent talk Sasa Prelovsek will present more details and recent results.

Statistical physics of isotropic-genesis nematic elastomers

BING-SUI LU

*Faculty of Mathematics and Physics
University of Ljubljana, Jadranska ulica 19, SI-1000 Ljubljana,
Slovenia
bing-sui.lu@fmf.uni-lj.si*

I will present an overview of nematic elastomers, which are nematic polymers that have been randomly, permanently cross-linked to form an equilibrium random solid network. I explain how such systems involve both annealed (thermal) as well as quenched (architectural) disorder. The distribution of thermal configurations of nematic polymers can be “memorized” by the elastomer at the instant of cross-linking, and I will show how a replica field theory formalism (inspired by Deam and Edwards) manages to capture this piece of physics. The random solid network restricts the motion of nematic polymer segments, and we address how this restriction modifies the liquid crystalline structure of nematic elastomers at high temperatures. In particular, we show that short-ranged oscillatory spatial correlations in the nematic alignment can emerge for a class of nematic elastomers at a sufficiently high density of cross-links.

References

- [1] B.-S. Lu, F. Ye, X. Xing, and P. M. Goldbart, *Physical Review Letters* **108** (2012) 257803.
- [2] B.-S. Lu, F. Ye, X. Xing, and P. M. Goldbart, *International Journal of Modern Physics B* **27** (2013) 1330012.

Wave dynamics in nonhermitian Photonics

KONSTANTINOS MAKRIS

Institute for Theoretical Physics

Technical University of Vienna (TU-Wien), A-1040, Vienna, Austria

Department of Electrical Engineering, Princeton University,

Princeton, 08544, NJ, USA

makris@princeton.edu

One of the frontiers of modern photonics is the engineering of the complex refractive index to create new composite systems with novel functionalities. In most systems optical loss has been always considered an obstacle. However, based on the recently introduced concept of parity-time (PT)-symmetry in the context of optical physics, such synthetic structures can utilize loss as an advantage and have been proven to be important for various nanophotonics applications. In this framework of open photonic systems, we also present a larger class of nonhermitian materials that are on average, lossy. These geometries exhibit transient power growth, the magnitude of which is related to the singular values of the non-normal propagator. The nontrivial wave dynamics in such nonhermitian optical structures can be understood in terms of the pseudospectra of the underlying non-normal operator. Active photonic systems that are on average lossy, are of current interest in nano-plasmonics.

References

- [1] C. M. Bender, *Reports on Progress in Physics* **70** (2007) 947-1018.
- [2] K. G. Makris, R. El-Ganainy, D. N. Christodoulides, and Z. H. Musslimani, *Physical Review Letters* **100** (2008) 103904.
- [3] K. G. Makris, R. El-Ganainy, D. N. Christodoulides, and Z. H. Musslimani, *Physical Review A* **81** (2010) 063807.
- [4] C. E. Rüter, K. G. Makris, R. El-Ganainy, D. N. Christodoulides, M. Segev, and D. Kip, *Nature Physics* **6** (2010) 192-195.
- [5] P. Ambichl, K. G. Makris, L. Ge, Y. Chong, A.D. Stone, and S. Rotter, *Physical Review X* **3** (2013) 041030.
- [6] K. G. Makris, L. Ge, and H. Türeci, *Physical Review X* (2014) accepted for publication.

Entropija in termoelektrični pojav v rutenatih

JERNEJ MRAVLJE

Odsek za teoretično fiziko

Institut Jozef Stefan, Jamova 38, SI-1000 Ljubljana, Slovenia

jernej.mravlje@ijs.si

V snoveh temperaturni gradienti povzročijo električno polje. Ta termoelektrični pojav nastane zaradi zlomitve simetrije delec-vrzeli in je uporaben v hlajenju in za koristno uporabo odvečne topote, npr. iz avtomobilskih izpuhov. Po drugi strani meritev razmerja med električnim poljem in termalnim gradientom (Seebeckov koeficient) vsebuje pomembno informacijo o snovi in je zanimiva v bazičnih raziskavah trdne snovi. V predavanju bom obravnaval zvezo med Seebeckovim koeficientom in entropijo. Kot primer bom uporabil snov z močnimi elektronskimi korelacijami Sr_2RuO_4 , ki jo bom opisal v teoriji dinamičnega povprečnega polja. Pokazal bom, da Seebeckov koeficient v ravnini ab tesno sledi rezultatu, ki sledi iz entropijskega razmisleka, Seebeckov koeficient v pravokotni ravnini c pa ne. To obnašanje razkriva poseben mehanizem, ki filtrira vrzeli in nastane kot posledica telesno centrirane rešetke.

Revealing active degrees of freedom, beating the entropy by hole filtering: lessons from thermopower in Sr₂RuO₄

JERNEJ MRAVLJE

*Department of theoretical physics
Jamova 38, SI-1000 Ljubljana, Slovenia
jernej.mravlje@ijs.si*

In materials, thermal gradients are accompanied by electrical fields. This 'thermoelectric effect' that comes from the particle-hole asymmetry can be used for solid-state refrigeration and waste heat recovery. On the fundamental side, the measurement of the ratio between the electric field and the thermal gradient (which is called Seebeck coefficient or thermopower) can be a revealing probe of the state of material. In this seminar, I will discuss the relation of thermopower to entropy. As an example, I will consider Sr₂RuO₄, a compound with intermediate strength of electronic correlations, which I will describe within the dynamical mean-field theory. I will argue that the in-plane thermopower reveals the active degrees of freedom which demonstrates potential of entropic interpretations of thermoelectric effect. On the other hand, the predicted c-axis thermopower is much higher and inconsistent with entropic consideration, a behavior that points to a peculiar hole-filtering mechanism.

Povezane strategije v evolucijskih igrah

MATJAŽ PERC

*Oddelek za fiziko, Fakulteta za naravoslovje in matematiko
Univerza v Mariboru, Koroška cesta 160, SI-2000 Maribor, Slovenia*

*CAMTP - Center za uporabno matematiko in teoretično fiziko
Univerza v Mariboru, Krekova 2, SI-2000 Maribor, Slovenia
matjaz_perc@uni-mb.si • www.matjazperc.com*

Kooperacija v človeških družbah je zelo pogosta in razširjena, in to nas pomembno ločuje od ostalih živalskih vrst. Začetki človeške kooperacije naj bi bili povezani s težavnostjo vzreje mladičev v zgodnji fazi razvoja naše vrste. Toda to že dolgo ni več problem. Vprašanje je, zakaj še vedno sodelujemo? Recipročnost je ena izmed ključnih predpostavk. Če je nekdo do nas prijazen, potem smo prijazni nazaj. Podobno, če smo izpostavljeni nepravičnosti in izkoriščevanju se maščujemo. Sodeč po teoriji močne recipročnosti naj bi bili tovrstna pozitivna in negativna recipročnost povezani. Toda je to res? Nedavni ekonomski eksperimenti zavračajo teorijo močne recipročnosti, in tudi vsakodnevne izkušnje govorijo, da ljudje bodisi primarno nagrajujemo lepo obnašanje ali pa kaznujemo grdo obnašanje, redko pa delamo z enako vnemo oboje. Pokazali bomo, kako lahko metode statistične fizike doprinesajo k razrešitvi nesoglasij med teorijo in prakso [1].

Reference

- [1] A. Szolnoki and M. Perc, Correlation of positive and negative reciprocity fails to confer an evolutionary advantage: Phase transitions to elementary strategies, *Physical Review X* **3** (2013) 041021.

Correlated strategies in evolutionary games

MATJAŽ PERC

*Department of Physics, Faculty of Natural Sciences and Mathematics
University of Maribor, Koroška cesta 160, SI-2000 Maribor, Slovenia*

*CAMTP - Center for Applied Mathematics and Theoretical Physics
University of Maribor, Krekova 2, SI-2000 Maribor, Slovenia
matjaz.perc@uni-mb.si • www.matjazperc.com*

Widespread cooperation among unrelated individuals distinguishes humans markedly from other species. The origins of our remarkable other-regarding abilities have been associated with rearing offspring that survived, which was a pressing challenge during the Paleolithic age that could not be met by individual efforts alone. But in the absence of such a challenge, what keeps us cooperating? Reciprocity is long considered an important piece of the puzzle. If someone is kind to us, we are kind in return. Similarly, if someone is unfair or exploitative, we tend to retaliate. And according to the strong reciprocity hypothesis, this positive and negative reciprocity are correlated to give us optimal evolutionary predispositions for the successful evolution of cooperation. But is this really true? Recent economic experiments reject the hypothesis, and everyday experiences leave us with the impression that people either tend to reward success or punish wrongdoing, but seldom will they do both. We show how methods of statistical physics contribute to resolving the disagreement between theory and observations [1].

References

- [1] A. Szolnoki and M. Perc, Correlation of positive and negative reciprocity fails to confer an evolutionary advantage: Phase transitions to elementary strategies, *Physical Review X* **3** (2013) 041021.

50 years of quarks

WILLIBALD PLESSAS

*Theoretical Physics, Institute of Physics
University of Graz, Universitätsplatz 5, A-8010 Graz, Austria
plessas@uni-graz.at*

Fifty years ago, in 1964, the notion of quarks was introduced [1]. Their interaction was formulated more than 40 years ago [2]. Still, the corresponding theory, quantum chromodynamics, cannot yet be solved comprehensively. Rather one has to resort to approximations, simulations, or models.

We discuss the development and present performance of the constituent-quark model for low-energy quantum chromodynamics [3,4], which gets by with a finite number of degrees of freedom contrary to the full quantum field theory. In particular, we address the spectroscopy of all known baryons, the structure of the nucleons exhibited under all (electromagnetic, weak, and gravitational) interactions, the electroweak structures of the other baryons with flavors up, down, and strange, as well as strong resonance decays. It turns out that by the improvements achieved over recent years a wide range of baryon properties and reactions can now be described by the relativistic constituent-quark model in rather good agreement with phenomenology and, where experimental data are missing, in reasonable congruence with results available from lattice quantum chromodynamics. The most important ingredients are relativistic invariance and a quark dynamics following from the spontaneous breaking of chiral symmetry in low-energy quantum chromodynamics. Challenging problems still exist with a proper description of resonance states and their couplings to decay channels.

References

- [1] M. Gell-Mann, *Phys. Lett.* **8** (1964) 214.
- [2] H. Fritzsch, M. Gell-Mann, and H. Leutwyler, *Phys. Lett. B* **47** (1973) 365.
- [3] L. Y. Glozman, W. Plessas, K. Varga, and R. F. Wagenbrunn, *Phys. Rev. D* **58** (1998) 094030.
- [4] J. P. Day, W. Plessas, and K. S. Choi, arXiv:1205.6918 [hep-ph].

Nered in antifragilnost v coulombskih tekočinah

RUDOLF PODGORNIK

*Oddelek za fiziko, Fakulteta za matematiko in fiziko, Univerza v
Ljubljana, and*

*Odsek za teoretično fiziko, J. Stefan Institut, SI-1000 Ljubljana,
Slovenija*

Kaj se zgodi z elektrostatskimi interakcijami, ?e je gostota naboja na makromolekulnih površinah podana z netermalizirano razurejeno porazdelitvijo? Predstavil bom formalni okvir [1], ki nam omogoča, da teoretično opišemo interakcije dolgega dosega v kontekstu omejenih coulombskih tekočin z razurejenimi izvori na robovih. Monopolaren in dipolaren nered vodita v tem kontekstu do bistveno modificiranih sil med mejnimi površinami [2], ki postanejo anomalno dolgega dosega tako v normalni kot tudi v lateralni smeri. Poleg tega se pojavijo tudi anomalni navori kot posledica razurejen porazdelitve nabojev. Posledice nereda ostajajo, tudi če so mejne površine v celoti vzeto elektroneutralne in lahko kot takšni razložijo celo vrsto parazitskih efektov, ki se jih da opaziti pri meritvah Casimirjeve sile. V primeru močno sklopljenih coulombskih tekočin z netermaliziranim neredom, razurejeni izvori lahko pripeljejo celo do pojava antifragilnosti v sistemu, ko več nereda na površinah vodi k zmanjšanju termičnega nereda.

Reference

- [1] Ali Naji, David S. Dean, Jalal Sarabadani, Ron R. Horgan, and Rudolf Podgornik, *Phys. Rev. Letts.* **104** (2010) 060601.
- [2] Jalal Sarabadani, Ali Naji, David S. Dean, Ron R. Horgan, and Rudolf Podgornik, *J. Chem. Phys.* **104** (2010) 1.

Disorder and antifragility in Coulomb fluids

RUDOLF PODGORNIK

*Department of Physics, Faculty of Mathematics and Physics,
University of Ljubljana, and
Department of Theoretical Physics, J. Stefan Institute, SI-1000
Ljubljana, Slovenia*

What happens to electrostatic interactions when charge density on macromolecular surfaces is characterized by quenched disorder distribution? Formal framework will be presented [1] that allows for the theoretical description of long-range electrostatic interactions in the presence of quenched disordered source fields on the boundaries. Monopolar and/or dipolar surface charge disorder, in the context of confined Coulomb fluids, gives rise to fundamentally modified interactions between planar bounding surfaces [2], characterized by anomalously long-ranged normal and lateral forces as well as torques. Disorder effects persist even in the case of overall charge-neutral surfaces and can help elucidate various parasitic effects known to exist in e.g. the Casimir interaction. In the case of strongly coupled Coulomb fluids with quenched disordered source fields a peculiar type of antifragility emerges where more imposed external source disorder actually diminishes the thermal disorder in the system, allowing the system to benefit from disorder.

References

- [1] Ali Naji, David S. Dean, Jalal Sarabadani, Ron R. Horgan, and Rudolf Podgornik, *Phys. Rev. Letts.* **104** (2010) 060601.
- [2] Jalal Sarabadani, Ali Naji, David S. Dean, Ron R. Horgan, and Rudolf Podgornik, *J. Chem. Phys.* **104** (2010) 1.

Močni razpadi mezonov $D_{s0}^*(2317)$ in $D_{sJ}(2460)$

ANITA PRAPOTNIK BRDNIK

*Univerza v Mariboru, Fakulteta za gradbeništvo,
Smetanova ul. 17, 2000 Maribor, Slovenija,
anita.prapotnik@um.si*

Leta 2003 je več kolaboracij opazilo ozki rezonanci $D_{s0}^*(2317)$ in $D_{sJ}(2460)$. Pridobljeni podatki nakazujejo, da gre za mezone pozitivne parnosti in $\bar{c}s$ strukture. Zaradi prenizke mase, omenjena stanja ne morejo razpasti v nižje ležeči D mezon z emitiranjem kaona, ampak se lahko emitirjo le pioni in fotoni. Ker pioni ne vsebujejo c ali s kvarka, močni razpadi so možni le preko kiralnih zank ali efektov izospinke kršitve, kot je naprimer mešanje $\eta - \pi_0$. Eksperimentalno so opazili tri takšne razpade: $D_{s0}^{*+}(2317) \rightarrow D_s^+ \pi_0$, $D_{sJ}(2460) \rightarrow D_s^{*+} \pi_0$ in $D_{sJ}(2460) \rightarrow D_s \pi^+ \pi^-$. Govorila bom o vplivu kiralnih zank na zgoraj naštete razpade z uporabo formalizma kiralne simetrije in limite težkih mezonov.

Reference

- [1] B. Aubert et al. (BABAR), *Phys. Rev.* **D74**, 031103 (2006).
- [2] B. Aubert et al. (BABAR), *Phys. Rev.* **D74**, 032007 (2006).
- [3] B. Aubert et al. (BABAR), *Phys. Rev. Lett.* **90**, 242001 (2006).
- [4] Y. Mikami et al. (Belle), *Phys. Rev. Lett.* **92**, 12002 (2004).
- [5] P. Colangelo, F. De Fazio, R. Ferrandes, *Mod. Phys. Lett.* **A19**, 2083 (2004).
- [6] W. Wei et al., *Phys. Rev.* **D73**, 034004 (2006).
- [7] J. Lu et al., *Phys. Rev.* **D73**, 054012 (2004).
- [8] W. Bardeen et al., *Phys. Rev.* **D68**, 054024 (2003).
- [9] I. W. Stewart, *Nucl. Phys.* **B529**, 62 (1998).
- [10] S. Fajfer, J. Kamenik, *Phys. Rev.* **D74**, 074023 (2006).
- [11] J. Zupan, *Eur. Phys. J.* **C25**, 233 (2002).
- [12] D. Bećirević, *Phys. Lett.* **B721**, 94 (2013).

Strong decay modes of $D_{s0}^*(2317)$ and $D_{sJ}(2460)$ mesons

ANITA PRAPOTNIK BRDNIK

*University of Maribor, Faculty of Civil Engineering,
Smetanova ul. 17, 2000 Maribor, Slovenia,
anita.prapotnik@um.si*

In 2003 several collaborations reported an observation of two narrow states: $D_{s0}^*(2317)$ and $D_{sJ}(2460)$. The data indicates that these are positive parity meson states with $\bar{c}s$ structure. Because of their low mass, $D_{s0}^*(2317)$ and $D_{sJ}(2460)$ can not decay into other D - meson states by emitting a kaon. Only emitting pions or photons is possible. As pions do not include strange quarks in their structure, the strong decay modes can only occur through chiral loops or isospin violating effects like $\eta - \pi$ mixing. Three such strong decay channels of $D_{s0}^*(2317)$ and $D_{sJ}(2460)$ were observed by experiment: $D_{s0}^{*+}(2317) \rightarrow D_s^+ \pi_0$, $D_{sJ}(2460) \rightarrow D_s^{*+} \pi_0$ and $D_{sJ}(2460) \rightarrow D_s \pi^+ \pi^-$. Here I will talk about the effects of chiral loops in these decay modes using the formalism of heavy meson chiral Lagrangians.

References

- [1] B. Aubert et al. (BABAR), *Phys. Rev.* **D74**, 031103 (2006).
- [2] B. Aubert et al. (BABAR), *Phys. Rev.* **D74**, 032007 (2006).
- [3] B. Aubert et al. (BABAR), *Phys. Rev. Lett.* **90**, 242001 (2006).
- [4] Y. Mikami et al. (Belle), *Phys. Rev. Lett.* **92**, 12002 (2004).
- [5] P. Colangelo, F. De Fazio, R. Ferrandes, *Mod. Phys. Lett.* **A19**, 2083 (2004).
- [6] W. Wei et al., *Phys. Rev.* **D73**, 034004 (2006).
- [7] J. Lu et al., *Phys. Rev.* **D73**, 054012 (2004).
- [8] W. Bardeen et al., *Phys. Rev.* **D68**, 054024 (2003).
- [9] I. W. Stewart, *Nucl. Phys.* **B529**, 62 (1998).
- [10] S. Fajfer, J. Kamenik, *Phys. Rev.* **D74**, 074023 (2006).
- [11] J. Zupan, *Eur. Phys. J.* **C25**, 233 (2002).
- [12] D. Bečirević, *Phys. Lett.* **B721**, 94 (2013).

Običajni in eksotični hadroni

SAŠA PRELOVŠEK KOMELJ

Fakulteta za Matematiko in Fiziko, Univerza v Ljubljani, Slovenija

Institut Jozef Stefan, Ljubljana, Slovenija

sasa.prelovsek@ijs.si

Kromodinamika na mreži predstavlja ne-perturbativni pristop za študij hadronskih lastnosti ab-initio. Predstavila bom dve mezonski stanji, ki se nahajata malce pod pragom in pri katerih simulacije prvič upoštevajo vpliv praga: gre za zanimivo stanje $X(3872)$ [1], ki je morda mezonska molekula $D^0\bar{D}^{0*}$, ter za skalarni mezon $D_{s0}^*(2317)$ [2,3]. Obravnavala bom tudi čudne [4] in čarobne [5] hadronske resonance, pri katerih smo poleg mase prvič določili tudi razpadno širino. Opisala bom zahtevno iskanje eksotičnih stanj s kvarkovsko strukturo $\bar{c}c\bar{d}u$ [6], ki so bila nedavo odkrita v eksperimentu.

Reference

- [1] S. Prelovsek, and L. Leskovec, *Phys. Rev. Lett.* **111**, 192001 (2013).
- [2] D. Mohler, C. B. Lang, L. Leskovec, S. Prelovsek, and R. M. Woloshyn, *Phys. Rev. Lett.* **111**, 222001 (2013).
- [3] C. B. Lang, L. Leskovec, D. Mohler, S. Prelovsek, and R. M. Woloshyn, *Phys. Rev.* **D90**, 034510 (2014).
- [4] S. Prelovsek, L. Leskovec, C. B. Lang, and D. Mohler, *Phys. Rev.* **D88**, 054508 (2013).
- [5] D. Mohler, S. Prelovsek, and R. M. Woloshyn, *Phys. Rev.* **D87**, 034501 (2013).
- [6] S. Prelovsek and L. Leskovec, *Phys. Lett.* **B727**, 172 (2013).

Conventional and exotic hadrons

SAŠA PRELOVŠEK KOMELJ

*Faculty of Mathematics and Physics, University of Ljubljana,
Jadranska 19, Slovenia
Jozef Stefan Institute, Jamova 39, Ljubljana, Slovenia
sasa.prelovsek@ijs.si*

QuantumChromoDynamics on the lattice presents a non-perturbative method to study the hadronic properties ab-initio. I will present the simulations of the mesonic states that lie slightly below thresholds, where the threshold effects are accounted for the first time. I will consider interesting state $X(3872)$ [1], which may be $D^0\bar{D}^{0*}$ molecule, and the scalar meson $D_{s0}^*(2317)$ [2,3]. The strange [4] and charmed [5] resonances will be considered, where the decay width is calculated for the first time. I will describe the challenging search for the exotic states with the quark content $\bar{c}c\bar{d}u$ [6], which have been recently discovered by several experiments.

References

- [1] S. Prelovsek, and L. Leskovec, *Phys. Rev. Lett.* **111**, 192001 (2013).
- [2] D. Mohler, C. B. Lang, L. Leskovec, S. Prelovsek, and R. M. Woloshyn, *Phys. Rev. Lett.* **111**, 222001 (2013).
- [3] C. B. Lang, L. Leskovec, D. Mohler, S. Prelovsek, and R. M. Woloshyn, *Phys. Rev.* **D90**, 034510 (2014).
- [4] S. Prelovsek, L. Leskovec, C. B. Lang, and D. Mohler, *Phys. Rev.* **D88**, 054508 (2013).
- [5] D. Mohler, S. Prelovsek, and R. M. Woloshyn, *Phys. Rev.* **D87**, 034501 (2013).
- [6] S. Prelovsek and L. Leskovec, *Phys. Lett.* **B727**, 172 (2013).

Točne rešitve kvantnih transportnih problemov daleč od ravnovesja in novi ohranitveni zakoni

TOMAŽ PROSEN

*Oddelok za fiziko, Fakulteta za matematiko in fiziko
Univerza v Ljubljani, Jadranska 19, Slovenija
tomaz.prosen@fmf.uni-lj.si • chaos.fmf.uni-lj.si*

Anizotropna Heisenbergova veriga spinov $1/2$ je osnovna paradigma sistema močno koreliranih elektronov v eni dimenziji (glej npr. [1]). Čeprav ravnovesne lastnosti modela dobro razumemo, so bile do nedavna odprta še zelo osnovna vprašanja v zvezi z neravnovesnimi, transportnimi lastnostmi, kot je npr. vprašanje [1,2], ali model dovoljuje balistični spinski transport pri končni temperaturi ali ne?

V predavanju bom na kratko orisal bistveni napredki na tem področju, ki ga je sprožilo odkritje točne rešitve kvantne master-enačbe za robno gnano Heisenbergovo verigo [3,4]. Stacionarna rešitev neravnovesne master enačbe namreč vodi do povsem novih kvazi-lokalnih ohranitvenih zakonov [3,7,10], ki posledično omogočajo strogo oceno balističnih transportnih koeficientov [6]. Takšen pristop ‘neravnovesne integrabilnosti’ deluje tudi za nekatere druge kvantne verige z močno interakcijo, kot je npr. Hubbardov model [8], ali Lai-Sutherlandov model verige spinov 1 [9].

Reference

- [1] X. Zotos in P. Prelovšek, “Transport in one dimensional quantum systems” poglavje v “Interacting Electrons in Low Dimensions”, serija “Physics and Chemistry of Materials with Low-Dimensional Structures”, Kluwer (2003).
- [2] J. Sirker, R. G. Pereira in I. Affleck, *Phys. Rev. Lett.* **103**, 216602 (2009); *Phys. Rev. B* **83**, 035115 (2011).
- [3] T. Prosen, *Phys. Rev. Lett.* **106**, 217206 (2011).
- [4] T. Prosen, *Phys. Rev. Lett.* **107**, 137201 (2011).
- [5] D. Karevski, V. Popkov in G. M. Schütz, *Phys. Rev. Lett.* **110**, 047201 (2013).
- [6] E. Ilievski in T. Prosen, *Commun. Math. Phys.* **318**, 809 (2013).
- [7] T. Prosen in E. Ilievski, *Phys. Rev. Lett.* **111**, 057203 (2013).
- [8] T. Prosen, *Phys. Rev. Lett.* **112**, 030603 (2014).
- [9] E. Ilievski in T. Prosen, *Nucl. Phys.* **B882**, 485 (2014).
- [10] T. Prosen, *Nucl. Phys.* **B886**, 1177 (2014).

Exact solutions of quantum transport problems far from equilibrium and new conservation laws

TOMAŽ PROSEN

Department of Physics, Faculty of Mathematics and Physics

University of Ljubljana, Jadranska 19, Slovenia

tomaz.prosen@fmf.uni-lj.si • chaos.fmf.uni-lj.si

Anisotropic Heisenberg chain of spins 1/2 is the key paradigm of strongly correlated electrons in one dimension (see e.g. [1]). While equilibrium properties of the model are relatively well understood, even very basic questions about its non equilibrium properties were still open until very recently. A prominent example is the question [1,2] whether the model exhibits ballistic spin transport at finite temperatures or not?

In my talk I will outline the progress on this topic which has been triggered by the discovery of exact solution of quantum master equation of the boundary driven Heisenberg chain [3,4]. The steady-state solution of non-equilibrium master equation leads to novel quasi-local conservation laws [3,7,10], which in turn lead to a derivation of rigorous strict lower bounds on ballistic transport coefficients [6]. Such an approach of ‘non-equilibrium integrability’ works also for some other strongly interacting quantum chains, for instance, for the Hubbard model [8] or Lai-Sutherland model of spins 1 [9].

References

- [1] X. Zotos in P. Prelovšek, “Transport in one dimensional quantum systems” poglavje v “Interacting Electrons in Low Dimensions”, serija “Physics and Chemistry of Materials with Low-Dimensional Structures”, Kluwer (2003).
- [2] J. Sirker, R. G. Pereira in I. Affleck, *Phys. Rev. Lett.* **103**, 216602 (2009); *Phys. Rev. B* **83**, 035115 (2011).
- [3] T. Prosen, *Phys. Rev. Lett.* **106**, 217206 (2011).
- [4] T. Prosen, *Phys. Rev. Lett.* **107**, 137201 (2011).
- [5] D. Karevski, V. Popkov in G. M. Schütz, *Phys. Rev. Lett.* **110**, 047201 (2013).
- [6] E. Ilievski in T. Prosen, *Commun. Math. Phys.* **318**, 809 (2013).
- [7] T. Prosen in E. Ilievski, *Phys. Rev. Lett.* **111**, 057203 (2013).
- [8] T. Prosen, *Phys. Rev. Lett.* **112**, 030603 (2014).
- [9] E. Ilievski in T. Prosen, *Nucl. Phys.* **B882**, 485 (2014).
- [10] T. Prosen, *Nucl. Phys.* **B886**, 1177 (2014).

Kvantni kaos v sistemih mešanega tipa

MARKO ROBNIK

*CAMTP - Center za uporabno matematiko in teoretično fiziko
Univerza v Mariboru, Krekova 2, SI-2000 Maribor, Slovenia
Robnik@uni-mb.si • www.camtp.uni-mb.si*

Najprej bom predstavil uvod v glavne pojme kvantnega kaosa, ali bolj splošno - valovnega kaosa, nato obravnaval sisteme mešanega tipa, to so generični Hamiltonovi sistemi, ki imajo klasično regularno dinamiko za določene začetne pogoje ter kaotično dinamiko za komplementarne začetne pogoje. Tipična struktura faznega prostora je kaotično morje, ki vsebuje otoke regularnosti in stabilnosti. Nato bom predstavil nekaj najnovejših izsledkov o kvantni lokalizaciji kaotičnih stanj.

Kvantna lokalizacija klasično kaotičnih lastnih stanj je eden najpomemnejših pojavov v kvantnem kaosu, poleg značilnega vedenja statističnih lastnosti energijskih spektrov. Kvantna lokalizacija nastopi, če je t.i. Heisenbergov čas t_H danega sistema krajši od vseh klasičnih transportnih časov v danem klasičnem sistemu, se pravi, kadar je klasični transport počasnejši od kvantne časovne resolucije evolucijskega operatorja. Heisenbergov čas t_H , kot pomembna karakteristika vsakega kvantnega sistema, je namreč enak razmerju Planckove konstante $2\pi\hbar$ ter srednjega razmika med energijskimi nivoji ΔE , $t_H = 2\pi\hbar/\Delta E$.

Pokazali bomo funkcionalno povezavo med stopnjo lokalizacije ter spektralno statistiko v avtonomnih (časovno neodvisnih) sistemih, v analogiji z brcanim rotorjem, ki pa je paradigma časovno periodičnih (Floquetovih) sistemov [7], ter pristop in metodo ilustrirali v primeru družine biljardov [8,9] v dinamičnem režimu med integrabilnostjo (krog) in polnem kaosom (kardioida), kjer bomo ekstrahirali kaotična stanja. Stopnjo lokalizacije določimo z dvema lokalizacijskima merama, z uporabo t.i. Poincaré-Husimijevih funkcij (ki so Gaussovo glajene Wignerjeve funkcije v Poincaré-Birkhoffovem faznem prostoru), ki so pozitivno definitne in jih lahko obravnavamo kot kvazi-verjetnostne gostote. Prva mera A je definirana s pomočjo informacijske entropije, druga (C), pa s pomočjo korelacij v faznem prostoru Poincaré-Husimijevih funkcij lastnih stanj. Presenetljivo in zelo zadovoljivo se izkaže, da sta obe meri linearno povezani in zato ekvivalentni.

Ena od glavnih manifestacij kaosa v kaotičnih stanjih v odsotnosti kvantne lokalizacije

je porazdelitev $P(S)$ razmikov S med sosednjimi energijskimi nivoji, ki se vede pri majhnih S kot linearna funkcija $P(S) \propto S$, in govorimo o linearinem odbijanju med sosednjimi nivoji, medtem ko imamo v integrabilnih sistemih Poissonovo statistiko (eksponentno funkcijo $P(S) = \exp(-S)$), kjer odbijanja med sosednjimi nivoji ni ($P(0) = 1 \neq 0$). V povsem kaotičnem režimu s kvantno lokalizacijo pa opazimo, da je $P(S)$ pri majhnih S potenčna funkcija $P(S) \propto S^\beta$, z $0 < \beta < 1$. Pokazali bomo, da obstaja funkcionalna odvisnost med mero lokalizacije A in eksponentom β , namreč da je β monotona funkcija A : pri močni lokalizaciji sta A in β majhna, pri šibki lokalizaciji (skoraj povsem razširjena kaotična stanja) pa sta A in β blizu 1.

Pristop ilustriramo v primeru kaotičnih stanj za zgoraj omenjene družine biljardov, kjer lahko ločimo regularna in kaotična stanja. Ta predstavitev sloni na naših najnovejših člankih [1,4,5,7].

Reference

- [1] B. Batistić and M. Robnik, *Phys.Rev. E* **88** (2013) 052913.
- [2] M.V. Berry and M. Robnik, *J. Phys.A: Math. & Gen.* **17** (1984) 2413.
- [3] B. Batistić and M. Robnik, *J. Phys. A: Math. & Theor.* **43** (2010) 215101.
- [4] B. Batistić and M. Robnik, *J. Phys. A: Math. & Theor.* **43** (2013) 315102.
- [5] B. Batistić, T. Manos and M. Robnik, *Europhys. Lett.* **102** (2013) 50008.
- [6] M. Robnik, *Nonlinear Phenomena in Complex Systems* (Minsk) **1** No. 1 (1998) 1-22.
- [7] T. Manos and M. Robnik, *Phys. Rev. E* **87** (2013) 062905.
- [8] M. Robnik, *J. Phys.A: Math. & Gen.* **16** (1983) 3971.
- [9] M. Robnik, *J. Phys.A: Math. & Gen.* **17** (1984) 1049.
- [10] T. Prosen and M. Robnik, *J. Phys.A: Math. & Gen.* **27** (1994) 8059.
- [11] T. Prosen and M. Robnik, *J. Phys.A: Math. & Gen.* **32** (1999) 1863.

Quantum chaos in the mixed-type systems

MARKO ROBNIK

CAMTP - Center for Applied Mathematics and Theoretical Physics

University of Maribor, Krekova 2, SI-2000 Maribor, Slovenia

Robnik@uni-mb.si • www.camtp.uni-mb.si

First I shall present an introduction to the main concepts of quantum chaos, or more generally - wave chaos, and then deal with the mixed-type systems, that are the generic Hamilton systems which have classical regular dynamics for some initial conditions and chaotic dynamics for the complementary initial conditions. The typical structure of the phase space is a chaotic sea containing islands of regularity and stability of motion. Then I will present some most recent results on quantum localization of chaotic eigenstates.

Quantum localization of classical chaotic eigenstates is one of the most important phenomena in quantum chaos, along with the characteristic behaviour of statistical properties of the energy spectra. Quantum localization sets in, if the Heisenberg time t_H of the given system is shorter than the classical transport times of the underlying classical system, i.e. when the classical transport is slower than the quantum time resolution of the evolution operator. The Heisenberg time t_H , as an important characterization of every quantum system, is namely equal to the ratio of the Planck constant $2\pi\hbar$ and the mean spacing between two nearest energy levels ΔE , $t_H = 2\pi\hbar/\Delta E$.

We shall show the functional dependence between the degree of localization and the spectral statistics in autonomous (time independent) systems, in analogy with the kicked rotator, which is the paradigm of the time periodic (Floquet) systems [7], and shall demonstrate the approach and the method in the case of a billiard family [8,9] in the dynamical regime between the integrability (circle) and full chaos (cardioid), where we shall extract the chaotic eigenstates. The degree of localization is determined by two localization measures, using the Poincaré Husimi functions (which are the Gaussian smoothed Wigner functions in the Poincaré Birkhoff phase space), which are positive definite and can be treated as quasi-probability densities. The first measure A is defined by means of the information entropy, whilst the second one, C , in terms of the correlations in the phase space of the Poincaré Husimi functions of the eigenstates. Surprisingly, and very satisfactory, the two measures

are linearly related and thus equivalent.

One of the main manifestations of chaos in chaotic eigenstates in absence of the quantum localization is the energy level spacing distribution $P(S)$ (of nearest neighbours), which at small S is linear $P(S) \propto S$, and we speak of the linear level repulsion, while in the integrable systems we have the Poisson statistics (exponential function $P(S) = \exp(-S)$), where there is no level repulsion ($P(0) = 1 \neq 0$). In fully chaotic regime with quantum localization we observe that $P(S)$ at small S is a power law $P(S) \propto S^\beta$, with $0 < \beta < 1$. We shall show that there is a functional dependence between the localization measure A and the exponent β , namely that β is a monotonic function of A : in the case of the strong localization are A and β small, while in the case of weak localization (almost extended chaotic states) A and β are close to 1.

We shall illustrate the approach in the model example of the above mentioned billiard family, where we can separate the regular and chaotic states. This presentation is based on our very recent papers [1,4,5,7].

References

- [1] B. Batistić and M. Robnik, *Phys.Rev. E* **88** (2013) 052913.
- [2] M.V. Berry and M. Robnik, *J. Phys.A: Math.& Gen.* **17** (1984) 2413.
- [3] B. Batistić and M. Robnik, *J. Phys. A: Math.& Theor.* **43** (2010) 215101.
- [4] B. Batistić and M. Robnik, *J. Phys. A: Math.& Theor.* **45** (2013) 315102.
- [5] B. Batistić, T. Manos and M. Robnik, *Europhys. Lett.* **102** (2013) 50008.
- [6] M. Robnik, *Nonlinear Phenomena in Complex Systems* (Minsk) **1** No. 1 (1998) 1-22.
- [7] T. Manos and M. Robnik, *Phys. Rev. E* **87** (2013) 062905.
- [8] M. Robnik, *J. Phys.A: Math.& Gen.* **16** (1983) 3971.
- [9] M. Robnik, *J. Phys.A: Math.& Gen.* **17** (1984) 1049.
- [10] T. Prosen and M. Robnik, *J. Phys.A: Math.& Gen.* **27** (1994) 8059.
- [11] T. Prosen and M. Robnik, *J. Phys.A: Math.& Gen.* **32** (1999) 1863.

Integrabilnost nekaterih sistemov NDE

VALERY G. ROMANOVSKI

CAMTP - Center za uporabno matematiko in teoretično fiziko

Univerza v Mariboru, Krekova 2, SI-2000 Maribor, Slovenia

Valery.Romanovsky@uni-mb.si • www.camtp.uni-mb.si

Obravnavamo problem iskanja sistemov z enim ali nekaj neodvisnimi prvimi integrali znotraj družin avtonomnih sistemov navadnih diferencialnih enačb odvisnih od parametrov. Metode konstruiranja prvih integralov ali dokazovanje njihovega obstoja v takšnih sistemih so prediskutirane. Med njimi je največ pozornosti usmerjene na Poincaré-Dulacovo metodo normalnih form, Darbouxjevo metodo in povezavo med integrabilnostjo in časovno reverzibilnostjo. Učinkovit računski pristop k problemu iskanja prvih integralov je predlagan. Nekateri primeri uporabe teorije in računskega pristopa so predstavljeni.

References

- [1] V. G. Romanovski, Y. Xia, X. Zhang, *Journal of Differential Equations* **257** (2014) 3079-3101.
- [2] Z. Hu, M. Aldazharova, T. M. Aldibekov and V. G. Romanovski, *Nonlinear Dynamics* **74** (2013) 1077-1092.
- [3] Z. Hu, M. Han and V.G. Romanovski, *Physica D* **265** (2013) 78-86.
- [4] V. G. Romanovski and M. Prešern, *Journal of Computational and Applied Mathematics* **236** (2011) 196-208.
- [5] V. G. Romanovski and D. S. Shafer *The Center and Cyclicity Problems: A Computational Algebra Approach*, Birkhäuser: Boston, 2009.

Integrability of some systems of ODEs

VALERY G. ROMANOVSKI

CAMTP - Center for Applied Mathematics and Theoretical Physics

University of Maribor, Krekova 2, SI-2000 Maribor, Slovenia

Valery.Romanovsky@uni-mb.si • www.camtp.uni-mb.si

The problem of finding systems with one or few independent first integrals inside families of autonomous systems of ODEs depending on parameters is discussed. Methods to construct first integrals or prove their existence in such systems are reviewed. Among them the main attention is paid to the Poincaré-Dulac normal form method, the Darboux method and the interconnection of integrability and time-reversibility. An efficient computational technique to the problem of finding first integrals is proposed. Some examples of application of the theory and the computational approach are presented.

References

- [1] V. G. Romanovski, Y. Xia, X. Zhang, *Journal of Differential Equations* **257** (2014) 3079-3101.
- [2] Z. Hu, M. Aldazharova, T. M. Aldibekov and V. G. Romanovski, *Nonlinear Dynamics* **74** (2013) 1077-1092.
- [3] Z. Hu, M. Han and V.G. Romanovski, *Physica D* **265** (2013) 78-86.
- [4] V. G. Romanovski and M. Prešern, *Journal of Computational and Applied Mathematics* **236** (2011) 196-208.
- [5] V. G. Romanovski and D. S. Shafer *The Center and Cyclicity Problems: A Computational Algebra Approach*, Birkhäuser: Boston, 2009.

Stability and periodic oscillations in the Moon-Rand System

DOUGLAS S. SHAFER

Mathematics Department

*The University of North Carolina at Charlotte, Charlotte, NC 28223
USA*

dsshafer@uncc.edu

The Moon-Rand system is a family of ordinary differential equations in three-dimensional Euclidean space developed to model control of flexible structures. Members of the family possess an equilibrium at the origin at which the linear part has one negative and two purely imaginary eigenvalues for all values of the parameters. We describe a complete stability analysis of the origin and provide a sharp upper bound on the number of isolated periodic orbits that can be made to bifurcate from the origin when the flow restricted to any so-called local “center manifold” is a focus. (Joint work with A. Mahdi and V. G. Romanovski.)

References

- [1] A. Mahdi, V. G. Romanovski, and D. S. Shafer, *Nonlinear Analysis: Real World Applications* **14** (2013) 294–313.
- [2] F. C. Moon and R. H. Rand, Jet Propulsion Laboratory Publication 85-29, vol. II, California Institute of Technology (2003) 329–342.
- [3] V. G. Romanovski and D. S. Shafer, *The Center and Cyclicity Problems: A Computational Algebra Approach*, Birkhäuser: Boston, 2009.

Stohastična simulacija prostorsko-časovne cirkadiane aktivnosti v suprakiazmatičnem jedru

Vito Šimonka, Maja Fras in Marko Gosak

*Fakulteta za naravoslovje in matematiko
Univerza v Mariboru, Koroška cesta 160, SI-2000 Maribor, Slovenia
vito.simonka@gmail.com*

Pri sesalcih dnevni ritem uravnava cirkadiani spodbujevalnik v suprakiazmatičnem jedru v hipotalamusu, ki organizem sinhronizira s periodičnimi okoljskimi pogoji, kot je dnevno-nočni svetlobni cikel. Najpomembnejša faktorja, ki zagotavlja 24-urni cirkadiani ritem, sta sklopitev nevronske aktivnosti z okoljem in medcelična komunikacija med nevroni [1]. Natančnost na časovni skali je dosežena kljub znatni ravni fluktuacij zaradi izredno majhnih koncentracij regulacijskih molekul. Da bi preverili vpliva molekularnega šuma in medcelične povezanosti na sinhronizacijo nevronske mreže, smo izdelali večcelični stohastični model suprakiazmatičnega jedra. Za opis medcelične komunikacije preko nevrotansmitterjev smo uporabili relativno realno skloplitveno shemo, ki temelji na eksperimentalnih ugotovitvah in izkazuje lastnosti mreže malega sveta [2,3]. Naši rezultati kažejo, da so nevronske mreže robustne na znotrajcelične fluktuacije in da jih lahko učinkovito vzbujamo s svetlobnimi cikli, kljub znatni meri stohastičnosti, pri čemer je koherenca oscilacij odvisna od narave medcelične komunikacije. Poleg tega smo pokazali, da vzorci nevronske aktivnosti oblikujejo lokalne sinhronizirane gruče, ki so bolj izrazite pri časovno kratkih fotoperiodah (zima) kot pri dolgih dnevih (poletje). S tem smo pokazali, da sezonske adaptacije cirkadianih ritmov ne vključujejo le reorganizacijo nevronske aktivnosti na časovni skali [3,4], temveč tudi na prostorski.

Reference

- [1] S. Yamaguchi, H. Isejima, T. Matsuo, R. Oktura, K. Yagita, M. Kabayashi, H. Okamura, *Science* **302** (2003) 1408-1412.
- [2] M. Hafner, H. Koeppl, D. Gonze, *PLoS computational Physics* **8** (2012) e1002419.
- [3] C. Bodenstein, M. Gosak, S. Schuster, M. Marhl, M. Perc, *PLoS computational biology* **8** (2012) e1002697.
- [4] J.H. Meijer, S. Michel, H.T. VanderLeest, J.H. Rohling, *European Journal of Neuroscience* **32** (2010) 2143-2151.

Stochastic simulation of the spatio-temporal circadian activity in the suprachiasmatic nucleus

Vito Šimonka, Maja Fras and Marko Gosak

*Fakulteta za naravoslovje in matematiko
Univerza v Mariboru, Koroška cesta 160, SI-2000 Maribor, Slovenia
vito.simonka@gmail.com*

In mammals, daily rhythms are driven by the circadian pacemaker in the suprachiasmatic nucleus (SCN) of the hypothalamus, which synchronizes an organism to periodically recurring environmental conditions, such as the solar day-night cycle. The coupling of the SCN neuronal activity with the environment and the intercellular communication between SCN neurons are the most important factors that ensure the precise 24h timing of the circadian rhythmicity, despite significant fluctuations in chemical activities due to very small numbers of regulatory molecules in individual cells [1]. We have developed a biochemically detailed stochastic multicellular model of the SCN in order to investigate the impact of molecular noise and intercellular connectivity on the circadian network synchronization. We have proposed a small-world coupling scheme for the intercellular communication via neurotransmitters [2,3]. Our findings reveal that the SCN network operates firmly despite significant levels of molecular noise, whereby the coherence of circadian dynamics relies significantly on the cell-to-cell interactions. We additionally show that the SCN neuronal activity patterns form localized clusters of synchronous behavior. This effect of spatial clustering is more pronounced in short days than in long photoperiods, thereby indicating that seasonal adaptation encompasses not only redistributions of the neuronal activity on the temporal level [3,4], but also on the spatial scale.

References

- [1] S. Yamaguchi, H. Isejima, T. Matsuo, R. Oktura, K. Yagita, M. Kabayashi, H. Okamura, *Science* **302** (2003) 1408-1412.
- [2] M. Hafner, H. Koeppl, D. Gonze, *PLoS computational Physics* **8** (2012) e1002419.
- [3] C. Bodenstein, M. Gosak, S. Schuster, M. Marhl, M. Perc, *PLoS computational biology* **8** (2012) e1002697.
- [4] J.H. Meijer, S. Michel, H.T. VanderLeest, J.H. Rohling, *European Journal of Neuroscience* **32** (2010) 2143-2151.

Tridimenzionalne faze tekočih kristalov iz paličastih molekul

NATAŠA VAUPOTIČ

*Fakulteta za naravoslovje in matematiko
Univerza v Mariboru, Koroška 160, SI-2000 Maribor, Slovenija
Institut "Jožef Stefan"
Jamova 39, SI-1000 Ljubljana, Slovenia
natasa.vaupotic@um.si • www.fnm.uni-mb.si*

Tekočekristalna faza je značilna za snovi, ki jih tvorijo močno anizotropne molekule. Največkrat je urejenost molekul v tekočekristalnih fazah eno ali dvodimenzionalna. Kljub izraziti anizotropni obliki molekul pa pogosto opazimo tekočekristalne faze s tridimenzionalno (3D) kubično ali tetragonalno strukturo [1]. 3D tekočekristalne faze pogosto sestojijo iz zlomljenih smektičnih plasti ali stolpcev molekul. Tvorba in vrsta 3D faz je močno odvisna od molekularnih parametrov, saj že npr. majhne spremembe v dolžini ogljikovodikovih repov lahko ključno vplivajo na fazno obnašanje sistema. Zelo pomemben vpliv imata tudi kiralnost in optična čistost.

Simetrijo kubične ali tetragonalne faze brez težav določimo z rentgenskim (rtg) sisanjem. Določitev notranje strukture kristalografske osnovne celice pa je veliko zahtevnejši problem. Z inverzno Fourierovo transformacijo rtg spektra lahko rekonstruiramo elektronsko gostoto v osnovni celici. Toda pri rtg sisanju dobimo le amplitudo strukturnega faktorja (kot koren iz intenzitete), izgubimo pa informacijo o fazi strukturnega faktorja. Problem lahko odpravimo tako, da skonstruiramo 3D model elektronske gostote in izračunamo Fourierovo transformacijo. Ustrezen model bo dal razmerja amplitud med vrhovi, ki so primerljiva z izmerjenimi, hkrati pa dobimo še faze vrhov. Elektronsko gostoto nato rekonstruiramo z uporabo eksperimentalno izmerjenih amplitud vrhov in teoretično dobljenih faz. Običajno je dovolj, če uporabimo le nekaj (dva do štiri) najvišjih vrhov, saj ostali ne vplivajo bistveno na elektronsko gostoto.

Predstavili bomo sistematično raziskavo vpliva kiralnosti, optične čistosti, doline ogljikovodikovih repov in njihove razvejanosti na 3D faze, ki jih tvorijo tekoči kristali iz paličastih molekul [2]. Poudarek bo na predstavitvi teoretičnih modelov in njihovi uporabi za identifikacijo struktur v kubičnih in tetragonalnih fazah.

Reference

- [1] C. Tschierske, *Angew. Chem. Int. Ed.* **52**, 8828 (2013).
- [2] M. Vogrin, N. Vaupotič, M. Wojcik, J. Mieczkowski, K. Madrak, D. Pociecha, E. Gorecka, *Phys. Chem. Chem. Phys.* **16**, 16067 (2014).

Three-dimensional phases of liquid crystals formed of rod-like molecules

NATAŠA VAUPOTIČ

*Faculty of Natural Sciences and Mathematics
University of Maribor, Koroška 160, SI-2000 Maribor, Slovenia
Jozef Stefan Institute
Jamova 39, SI-1000 Ljubljana, Slovenia
natasa.vaupotic@um.si • www.fnm.uni-mb.si*

Thermotropic liquid crystalline cubic and tetragonal phases are three-dimensionally (3D) ordered, highly symmetric structures, which can be formed also by strongly anisotropic mesogenic molecules. They are relatively common among the rod-like, polycatenar, polyhydroxy and dendritic molecules [1]. The 3D density modulated phases are usually made from strongly distorted smectic layers or columns, depending on the type of constituent molecules. Formation of the 3D phases is very sensitive to molecular factors, e.g. even small changes in the length of terminal chains might have a crucial effect on the behavior of the system. Chirality and optical purity play an important role, as well.

Identification of the symmetry of cubic or tetragonal phases is usually straightforward from the X-ray diffraction pattern. On the other hand, revealing the internal structure of crystallographic unit cell is an intriguing task. It could be done by the reconstruction of the electron density map through the reverse Fourier transform of the X-ray pattern, however in a standard X-ray experiment only information related to the amplitude of structure factor (related to the square root of signal intensity) is available, while the information about the structure factor phase is lost. Moreover, for cubic phases multiplicity of signals is high and peaks belonging to the same multiplicity set might have different phases. One way to solve this problem is to construct a model of a 3D electron density distribution and calculate its Fourier transform. An appropriate model should yield the signal amplitude ratios close to the experimental ones. Finally, the electron density map is reconstructed using experimental amplitudes and theoretically obtained phases. Usually it is enough to use only few signals of the highest intensity; the other peaks do not affect the electron density significantly.

In the talk we will report on detailed X-ray diffraction studies that enabled a reconstruction of the electron density distribution in cubic as well as, for the first time, in tetragonal LC phases. Systematic studies of the effect of chirality, optical purity, length of terminal chains and their branching on the 3D packing of rod-like mesogenic molecules were conducted [2].

References

- [1] C. Tschierske, *Angew. Chem. Int. Ed.* **52**, 8828 (2013).
- [2] M. Vogrin, N. Vaupotič, M. Wojcik, J. Mieczkowski, K. Madrak, D. Pociecha, E. Gorecka, *Phys. Chem. Chem. Phys.* **16**, 16067 (2014).

1. Koncert / 1st Concert

Nikolaj Sajko, cello

Miha Haas, piano

Program:

Ludwig van Beethoven: Sonata for piano and cello in A-major, Op. 69

- . I. Allegro, ma non tanto
- . II. Scherzo. Allegro molto
- . III. Adagio cantabile - Allegro vivace

Blaž Pucihar: Summer Sonata

Astor Piazzolla: Le Grand Tango

Nikolaj Sajko graduated in cello performance from the Academy of Music in Ljubljana in the class of Ciril Škerjanec and completed his Master of Science at the same institution, researching the articulation and phrasing of the orchestral playing. As a soloist he perfected his performance further at postgraduate studies at the Anton Bruckner Private University in Linz. During his studies he regularly took part in several international seminars with prominent professors such as I. Gavriš. E. Schoenfeld, A. Noras, C. Onczay and T. Khne. Nikolaj Sajko was a member of the Gustav Mahler youth orchestra which operates under the leadership of Claudio Abbado. Besides that he was a member of the World Youth Orchestra. As a soloist Nikolaj performed with Maribor Philharmonic Orchestra, chamber orchestras Gaudeamus and Amadeus and with Maribor Youth Orchestra. He has performed in several solo recitals, chamber concerts and successful musical projects on the national as well as the international level. He was awarded the Klasinc diploma and Antonio Tarsia prize; other important achievements come from national and international competitions for young musicians in Gorizia, Lienz, Zagreb and Ljubljana. At present he is assistant solo cellist in the Symphony Orchestra of Slovene National Theatre in Maribor and a professor at the Conservatory for Music and Ballet in Maribor. In January this year Nikolaj released his first duo CD, along with pianist Miha Haas, with works for cello and piano by Edvard Grieg, Risto Savin, Bohuslav Martin and Blaž Pucihar. Since 2012 Nikolaj is artistic leader of chamber music series Carpe artem, which is co-organized by Musicians' Society Amadeus and Slovene National Theatre Maribor.

Miha Haas graduated in piano performance from the Academy of Music in Ljubljana in the class of Dubravka Tomšič-Srebotnjak and finished his post-graduation at The Royal Music Conservatory in Brussels in the class of Aleksandar Madžar. He took part at the international music seminars for piano and chamber music and worked with I. Lasko, J. Siirala, P. Paleczny, J. M. Huizing, M. Voskresenski, P. Kuijken and S. Picard. As a soloist he played with the Slovenian Philharmonic

Orchestra, The RTV Slovenia Symphony Orchestra, The Symphonic Orchestra of the Academy of Music Ljubljana, The Symphonic Orchestra of Ljubljana Secondary School of Music, Celje String Orchestra, The Wind Orchestra of the Slovenian Army, The Wind Orchestra Ravne, and Festival Bled Orchestra. Moreover, his musical activities include also solo recitals, chamber music concerts especially in piano duos and in duos with the violin or cello. He participated in premier performances of 8 compositions by Slovenian composers and he made several recordings for the national radio and television. He was awarded the Prešeren prize of the Academy of Music Ljubljana and the Prešeren prize of the University of Ljubljana. Together with the pianist Tadej Horvat with whom he collaborates in a piano duo he won the second prize and the special prize for the best performance of the piece by E. Grieg at the Concours Grieg 2009 in Oslo. At present he works at the Academy of Music in Ljubljana.

Nikolaj Sajko je diplomiral iz violončela na Akademiji za glasbo v Ljubljani v razredu profesorja Cirila Škerjanca. Na ljubljanski Akademiji za glasbo je tudi končal znanstveni magistrski študij na temo artikulacije in faziranja v orkestrski igri, podiplomsko pa se je v solistični igri dodatno izpopolnjeval na Univerzi Antona Brucknerja v Linzu in Univerzi za umetnost v Gradcu. Med študijem se je udeleževal mednarodnih seminarjev pri priznanih profesorjih I. Gavrišu, E. Schoenfeld, A. Norasu, C. Onczayu in T. Khneju. Bil je član mednarodnega mladinskega orkestra Gustav Mahler, ki deluje pod umetniškim vodstvom Claudia Abbada, in član Svetovnega orkestra glasbene mladine. Kot solist je nastopil z orkestrom Mariborske filharmonije, s komornim orkestrom Gaudeamus, s komornim orkestrom Amadeus in z Mariborskим mladinskim godalnim orkestrom. V dosedanjem glasbenem delovanju je izvedel več solističnih recitalov, komornih koncertov ter odmevnih glasbenih projektov tako na državni kot na mednarodni ravni. Je dobitnik Klasinčeve diplome in nagrade Antonia Tarsie, pomembne uvrstitve pa je dosegel tudi na državnih in mednarodnih tekmovanjih mladih glasbenikov v Gorici, Lieznu, Zagrebu in Ljubljani. Je namestnik soločelista v Simfoničnem orkestru SNG Maribor ter poučuje na Konservatoriju za glasbo in balet v Mariboru. Leta 2011 je skupaj s pianistom Miho Haasom izdal svojo prvo CD-ploščo z deli za violončelo in klavir Edvarda Griega, Rista Savina, Bohuslava Martinja in Blaža Puciharja. Od leta 2012 je umetniški vodja cikla koncertov komorne glasbe Carpe artem, ki jih organizirata Društvo za komorno glasbo Amadeus in Slovensko narodno gledališče Maribor.

Miha Haas je diplomiral iz klavirja na Akademiji za glasbo v Ljubljani, v razredu profesorce Dubravke Tomšič-Srebotnjak, podiplomski študij pa dokončal na Kraljevem konservatoriju v Bruslju, v razredu profesorja Aleksandra Madžarja. Izpopolnjeval se je na mednarodnih seminarjih za solistično in komorno igro pri I. Lasku, J. Siirali, P. Palecznyju, J. M. Huizingu, M. Voskresenskem, P. Kuijkenu in S. Picardu. Kot solist je nastopil z Orkestrom Slovenske filharmonije, z Orkestrom RTV Slovenija, s Simfoničnim orkestrom Akademije za glasbo v Ljubljani, z Orkestrom Srednje glasbene in baletne šole Ljubljana, s Celjskim godalnim orkestrom, s Pihalnim orkestrom Slovenske vojske, s Pihalnim orkestrom Železarjev Ravne in z

Orkestrom festivala Bled. Njegovo glasbeno delovanje ob koncertih z orkestri zajema še izvedbe solističnih recitalov, nastopov z različnimi komornimi zasedbami, predvsem v klavirskem duu ter v duih z violino in čelom. Sodeloval je pri krstnih izvedbah osmih novitet slovenskih avtorjev ter snemal za arhiv televizije in radia Slovenija. Prejel je škerjančevo nagrado Srednje glasbene in baletne šole Ljubljana, Prešernovo nagrado Akademije za glasbo ter Prešernovo nagrado Univerze v Ljubljani. S pianistom Tadejem Horvatom, s katerim deluje v klavirskem duu, sta osvojila drugo nagrado ter posebno nagrado za najboljšo interpretacijo skladbe E. Griega na tekmovanju Concours Grieg 2009 v Oslu. Danes je zaposlen na Akademiji za glasbo v Ljubljani.

2. Koncert / 2nd Concert

Nejc Kamplet

Program:

1. J. S. Bach (1685-1750) Prelude and fugue in E flat BWV 876
2. . Frederic Chopin (1810-1849) Heroic Polonaise Op. 53 in A flat major
3. Ludwig van Beethoven (1770-1827) Sonata No. 21 in C major, Op. 53, Waldstein
 - . allegro con brio
 - . adagio molto
 - . rondo, allegretto moderato - prestissimo
4. F. Liszt (1811-1886) Mephisto Waltz No. 1, S. 514
5. Robert Kamplet (1971) Meditation
6. F. Chopin (1810-1849) Sonata No. 3 in B minor, Op. 58 - allegro maestoso
7. A. Volodos (1972) Concert Paraphrase on Mozart's Turkish March



Nejc Kamplet, coming from Slovenia is son of the composer and the flautists and is learning the piano since the age of five. He graduated from music in the Conservatory in Maribor with Distinction (class of Sasa G. Donaldson). He is now a student of a high school in Maribor (Slovenia) and at the same time he is studying at the University of Music and performing Arts Graz (Kunstuniversitat Graz, Austria) in the class of Dr. Suzana Niederdorfer. He has received many awards and first places at the national and international competitions. The major awards during the recent time are: golden plaques and special awards for the best performances of pieces by Slovenian composers in national competition TEMSIG in the years 2007, 2010 and 2013, first prize at the international competition Zlatko Grgosevic in Zagreb, Croatia (2008), first prize and special prize for the best performance of classical sonatas at the competition of Ars Nova in Trieste (2008) and WPW Panmusica in Vienna (2009) (second prize). In the year 2013 he won the first prize (absolute points) in competition T. Holmar in Malborghetto (Italy) and second prize at the piano competition Giovanni Musicisti in Trevisio (Italy). In the August 2014, he performed in intense competition in semi-finals at the 14th Ettlingen International Competition. He also played in the Infinitum piano trio who won the first prize at the national competition TEMSIG (2011) for chamber music. Nejc Kamplet regularly attends master classes in Slovenia, Germany, Croatia, Liechtenstein, Estonia, Slovakia, etc. He is participating in many music festivals in Slovenia (Festival Lent, Festival Maribor, etc.) and he also played many piano concertos with many orchestras. Most noticeable are: Mozart Concertos K41, K482, K365 (2007, 2010, 2011), Beethoven first Piano Concerto in C major (2012), E. Grieg first Piano Concerto in a minor (2013) and in January 2014 he played his fathers R. Kamplet first piano concerto with SNG (Maribor symphony Orchestra) with conductor Tae Jung Lee. He also participated with many other renowned conductors (Benjamin Pionnier, Slavko Magdic, Ziva P. Persuh, etc.). In 2010 he received the prize Dr. R. Klasinc by the Conservatory of Music and Ballet Maribor for the best students.

Kazalo / Contents

Organizatorji / Organizing Committee	1
Predgovor	2
Foreword	3
Seznam udeležencev / Participants list	4
Urnik / schedule	7
Povzetki / Abstracts	11
1. Koncert / 1st Concert	66
2. Koncert / 2nd Concert	69